

ENDEMIC PLANTS OF THE CENTRAL GRASSLAND OF NORTH AMERICA: DISTRIBUTION, ECOLOGY, AND CONSERVATION STATUS

James H. Locklear

Lauritzen Gardens

100 Bancroft Street

Omaha, Nebraska 68108, U.S.A.

j.locklear@omahabotanicalgardens.org

ABSTRACT

This paper enumerates the endemic plants of the Central Grassland of North America. The Central Grassland encompasses the full extent of the tallgrass, mixed-grass, and shortgrass prairie ecological systems of North America plus floristically related plant communities that adjoin and/or interdigitate with the midcontinental grasslands including savanna-open woodland systems, shrub-steppe, and rock outcrop communities. There are 382 plant taxa endemic to the Central Grassland, 300 endemic species (eight of which have multiple subspecific taxa endemic to the region) and 72 endemic subspecies/varieties of more widely distributed species. Nine regional concentrations of endemic taxa were identified and are described as centers of endemism for the Central Grassland: Arkansas Valley Barrens, Edwards Plateau, Llano Estacado Escarpments, Llano Uplift, Mescalero-Monahans Dunes, Niobrara-Platte Tablelands, Raton Tablelands, Red Bed Plains, and Reverchon Rocklands. In addition to hosting localized endemics, these areas are typically enriched with more widely-distributed Central Grassland endemics as well as peripheral or disjunct occurrences of locally-rare taxa, making them regions of high floristic diversity for the Central Grassland. Most of the endemics (299 or 78%) are habitat specialists, associated with rock outcrop, sand, hydric, or riparian habitats. There is a strong correlation between geology and endemism in the Central Grassland, with 59% of the endemics (225 taxa) associated with rock outcrop habitat. Of the 382 Central Grassland endemics, 124 or 33% are of conservation concern (NatureServe ranking of G1/T1 to G3/T3). Of these at-risk taxa, 78 or 63% are primarily associated with one of the centers of endemism identified in the study. It is hoped these findings will be useful in focusing conservation action on the habitats, ecological associations, and regions of the Central Grassland that host the highest concentrations of unique and at-risk plant species and associated biological diversity.

KEY WORDS: Central Grassland, conservation, endemism, floristic diversity, geobotany

RESUMEN

En este artículo se enumeran las plantas endémicas de la Pradera Central de Norteamérica. La Pradera Central abarca totalidad de los sistemas ecológicos de praderas de gramíneas altas, mezcladas, y bajas de Norteamérica junto con comunidades de plantas relacionadas florísticamente que se añaden y/o se entremezclan con las praderas continentales que incluyen sistemas de sabana-bosque abierto, arbusto-estepa, y comunidades de afloramientos rocosos. Hay 382 taxa endémicos de la Pradera Central, 300 especies endémicas (ocho de las cuales tienen múltiples taxa subspecíficos endémicos de la región) y 72 subspecies/variedades endémicas de especies de distribución más amplia. Se identificaron nueve concentraciones regionales de taxa endémicos y se describen como centros de endemismos de la Pradera Central: Arkansas Valley Barrens, Edwards Plateau, Llano Estacado Escarpments, Llano Uplift, Mescalero-Monahans Dunes, Niobrara-Platte Tablelands, Raton Tablelands, Red Bed Plains, y Reverchon Rocklands. Además de albergar endemismos localizados, estas áreas están típicamente enriquecidas con endemismos de la Pradera Central de distribución más amplia, así como ocurrencias periféricas o disjuntas de taxa localmente raros, que hacen de ellas regiones de alta diversidad florística para la Pradera Central. La mayoría de los endemismos (299 o 78%) están especializados en un hábitat, asociados con afloramientos rocosos, arena, agua, o hábitats riparios. Hay una correlación fuerte entre geología y endemismo en la Pradera Central, con 59% de los endemismos (225 taxa) asociados con afloramientos rocosos. De los 382 endemismos de la Pradera Central, 124 o 33% necesitan conservación (ranking de NatureServe de G1/T1 a G3/T3). De estos taxa en riesgo, 78 o 63% están asociados primariamente con uno de los centros de endemismo identificados en el estudio. Se espera que estos hallazgos sean útiles para establecer las acciones de conservación en los hábitats, asociaciones ecológicas, y regiones de la Pradera Central que alberga las mayores concentraciones de especies únicas y amenazadas y la diversidad biológica asociada.

INTRODUCTION

“The prairies have very few endemic plants.”—*P.A. Rydberg* (1931)

“The relative shortage of endemics in the grassland vegetation is a significant fact.”—*P.V. Wells* (1970)

“It is a striking fact that the Central Plains flora is not rich in unique taxa.”—*D.I. Axelrod* (1985)

Knowledge of endemism in the flora of a region is an important element in understanding the biological

diversity and distinctiveness of the region and is helpful in prioritizing areas for conservation action. An endemic species or subspecific taxon is one whose geographic distribution is limited to some particular region. Depending on the geographic scale of the area under consideration, an endemic may have a relatively large area of distribution and may even be abundant within its range. Rare or narrow endemics are those having small ranges, high habitat specialization, low local abundance, or a combination of these attributes (Kruckeberg & Rabinowitz 1985). Generally, the narrower the distribution of a species, the greater the conservation concern.

Assessment of endemism requires a well-developed knowledge of a region's flora. Of the relatively few published enumerations of endemic plants for North America, most are limited to a particular state. Examples include Arkansas (Gentry et al. 2013), Colorado (Weber & Wittmann 1992), New Mexico (Spellenberg 1993; Sivinski & Knight 1996), Texas (Carr 2009), Utah (Shultz 1993), and Virginia (Weakley et al. 2012). California has the most extensive plant endemism literature of any state (Stebbins & Major 1965; Raven & Axelrod 1978; Smith & Sawyer 1988), including recent analyses of the biogeography of California endemic plants and the implications for conservation planning (Thorne et al. 2009; Kraft et al. 2010). Enumeration of endemic plants for regions that encompass more than one state or province are even fewer in number, but include the Atlantic and Gulf Coastal Plain (Sorrie & Weakley 2001; MacRoberts et al. 2002), the Interior Highlands (Zoller et al. 2005), the Four Corners region (Arizona, Colorado, New Mexico and Utah) of the southwest (Heil et al. 2013), and the California Floristic Province (Burge et al. 2016).

This paper enumerates the endemic plants of the grasslands and related plant communities of central North America. It is focused on the three major midcontinental grassland systems—tallgrass prairie, mixed-grass prairie and shortgrass prairie—which R. Anderson (2006) referred to collectively as the Central Grassland of North America. It roughly corresponds to the North American Prairies floristic province of Takhtajan (1986).

The study also includes floristically-related plant communities that adjoin and/or interdigitate with the midcontinental grasslands including savanna-open woodland systems, shrub-steppe, and rock outcrop communities. While not grassland or prairie vegetation per se, these associations possess a significant component of grasses, graminoids, and forbs characteristic of the regional grassland flora. This approach parallels that taken by Noss (2013) in his comprehensive survey of the grasslands of the southern United States in which he defined grassland as “any natural community or ecosystem in which the herbaceous layer is dominated by grasses, other graminoid (grass-like) plants such as sedges, and associated forbs (other herbaceous plants).” Such plant communities are here recognized collectively as “campestrian,” a term used in biogeography to designate regions dominated by grassland vegetation (Cooper 1859; Mohr 1901; Turner 1959; Armstrong 1972). A synopsis of the NatureServe ecological systems encompassed by this study is presented in Table 1.

Previous Assessments of Endemism in the Central Grassland

Rydberg was the first to catalog the flora of the midcontinental grasslands, publishing *Flora of the Rocky Mountains and Adjacent Plains* in 1917 and *Flora of the Prairies and Plains of Central North America* in 1932. He also was first to comment on endemism in the flora, making the statement above in “A Short Phytogeography of the Prairies and Great Plains of Central North America.” In this paper, Rydberg (1931) used “the prairies” to denote tallgrass prairie (“tall grass lands”) and “the plains” to denote grasslands to the west (“short grass lands”). He provided no list or number of endemics or even examples of species he considered endemic, so it is unclear whether he was referring to all of the grasslands of central North America or only the tallgrass prairie of the east. He stated later in the paper that “about 25 species” are endemic to the “Staked Plains”—his term for the region of the western Great Plains south of the divide between the Arkansas and Platte rivers—but again provided no specific data.

The next published assessment of endemism in the flora of the region came nearly 40 years later in Wells' 1970 paper, “Historical Factors Controlling Vegetation Patterns and Floristic Distributions in the Central Plains Region of North America.” Using Rydberg's (1932) *Flora of the Prairies and Plains of Central North America* for his analysis, Wells estimated that about five percent of the flora of the grassland vegetation of the Interior Plains was endemic. Rydberg's treatment included 3,988 species, which Wells distilled down to 954

TABLE 1. Synopsis of NatureServe ecological systems encompassed within study region.

Prairies	Shrubland
Central Mixedgrass Prairie	Edwards Plateau Limestone Shrubland
Central Tallgrass Prairie	Llano Estacado Caprock Escarpment and Breaks Shrubland and Steppe
North-Central Interior Sand and Gravel Tallgrass Prairie	Northwestern Great Plains Shrubland
Northern Great Plains Fescue-Mixed Grass Prairie	Western Great Plains Mesquite Scrub Woodland and Shrubland
Northwestern Great Plains Mixedgrass Prairie	Western Great Plains Sandhill Steppe
Southeastern Great Plains Tallgrass Prairie	Barrens, Glades, and Outcrops
Texas Blackland Tallgrass Prairie	Central Interior Highlands Calcareous Glade and Barrens
Western Great Plains Foothill and Piedmont Grassland	Edwards Plateau Carbonate Glade and Barrens
Western Great Plains Sand Prairie	Llano Uplift Acidic Forest, Woodland and Glade
Western Great Plains Shortgrass Prairie	Northwestern Great Plains Canyon
Western Great Plains Tallgrass Prairie	Southeastern Great Plains Cliff
Savanna and Woodland	Southwestern Great Plains Canyon
Cross Timbers Oak Forest and Woodland	Western Great Plains Badlands
Eastern Great Plains Tallgrass Aspen Parkland	Western Great Plains Cliff and Outcrop
Edwards Plateau Limestone Savanna and Woodland	Herbaceous Wetland
North-Central Interior Oak Savanna	Eastern Great Plains Wet Prairie, Meadow and Marsh
North-Central Oak Barrens	Edwards Plateau Playa
Northwestern Great Plains Aspen Forest and Parkland	Edwards Plateau Riparian
Northwestern Great Plains-Black Hills Ponderosa Pine Woodland and Savanna	Great Plains Prairie Pothole
Ozark Prairie and Woodland	North-Central Interior Freshwater Marsh
Southern Rocky Mountain Juniper Woodland and Savanna	Western Great Plains Closed Depression Wetland and Playa
Southern Rocky Mountain Pinyon-Juniper Woodland	Western Great Plains Open Freshwater Depression Wetland
Southern Rocky Mountain Ponderosa Pine Savanna	Western Great Plains Riparian
Western Great Plains Dry Bur Oak Forest and Woodland	Western Great Plains Saline Depression Wetland

species that occurred naturally “in grassland vegetation on the plains.” Five percent of this more restricted flora would translate into roughly 50 endemic species.

On a more limited scale, Bare and McGregor (1970) mapped and analyzed the distribution patterns of 479 vascular plants species native to Kansas and determined that only six species in the Kansas flora were endemic to the Great Plains. Five of these are recognized herein as endemic to the Central Grassland: *Aster fendleri* (= *Symphotrichum fendleri*), *Juncus brachyphyllus*, *Oenothera fremontii* (= *O. macrocarpa* subsp. *fremontii*), *Phlox oklahomensis*, and *Scutellaria resinosa*.

The most recent assessment of endemism in the flora of the region appears in “Rise of the Grassland Biome, Central North America” by Axelrod (1985). Axelrod’s analysis benefited from the detailed distribution maps published in *Atlas of the Flora of the Great Plains* (GPFA 1977), by which he determined that 68 taxa were endemic to “the central grassland region” and 322 “almost endemic, but with some sites in bordering areas.” Axelrod limited his analysis to herbaceous and perennial taxa and excluded aquatic, near-aquatic, and riparian families such as Salicaceae and Potamogetonaceae.

In his overview of phytogeography for *Flora of North America* Volume 1, Thorne (1993) echoed Rydberg, Wells, and Axelrod in stating, “endemism is rather limited in the Prairies Province,” noting “no endemic families, few if any endemic genera, and perhaps fewer than 50 endemic species.”

Resources for a New Assessment

Publication of *Flora of the Great Plains* (GPFA 1986) was a crucial milestone in documenting the flora of the midcontinental grasslands and represented the culmination of decades of field and herbarium work by a team of researchers located at colleges and universities throughout the Great Plains. Even so, contributor Robert Kaul noted at the time of publication, “Basic floristic information is lacking for large parts of our area.” Today, students of the flora of the region have many more resources at hand to aid their work, notably updated state floristic references for Colorado (Ackerfield 2015; Weber and Wittmann 2012), Missouri (Yatskievych 1999, 2006, 2013), Montana (Lesica 2012), Nebraska (Kaul et al. 2011), Texas (Diggs et al. 1999; Turner et al. 2003),

and Wyoming (Dorn 2001). Additionally, smaller scale floristic inventories and vegetation studies have been published for special areas within the Plains, including several of the U.S.D.A. Forest Service National Grassland units (Hazlett 1998, 2004; Hazlett et al. 2009; Kuhn et al. 2011). These resources, coupled with recent phylogenetic studies of groups with Plains taxa, many in preparation for the *Flora of North America* (FNAEC 1993–), provide data for a new assessment of endemism in the flora of the Central Grassland.

STUDY REGION

The core of the study region is the Central Grassland of North America as delineated by R. Anderson (2006) and encompasses the full extent of the tallgrass, mixed-grass, and shortgrass prairies of the midcontinent. This descriptive name is abbreviated hereafter as “Central Grassland” and endemics of the region as “CG endemics.” Figure 1 shows the geographic extent of the study region.

The study region encompasses but is not limited to the Great Plains of North America. Fenneman’s (1931) physiographic definition of the Great Plains is widely used in ecological and biogeographic studies but the boundaries of Fenneman’s Great Plains physiographic province excludes most of the tallgrass prairie and much of the southern mixed-grass prairie. The study region is also more expansive than that considered in *Atlas of the Flora of the Great Plains* (GPFA 1977) and *Flora of the Great Plains* (GPFA 1986), which exclude significant reaches of the midcontinental grasslands that lie to the east, south and north (see Figure 1). The use of the term “Great Plains” or “Plains” is limited hereafter to the physiographic province as defined by Fenneman.

Thus circumscribed, the study region takes in all or part of 17 states—Arkansas, Colorado, Illinois, Indiana, Iowa, Kansas, Minnesota, Missouri, Montana, Nebraska, New Mexico, North Dakota, Oklahoma, South Dakota, Texas, Wisconsin, and Wyoming—plus the Canadian provinces of Alberta, Manitoba, and Saskatchewan. Finer-scale, ecosystem-level boundaries are delineated below. A synopsis of NatureServe ecological systems encompassed by this study is presented in Table 1.

Northern Extent

The study region extends north of the United States to encompass the prairie grasslands of Canada, including the Aspen Parkland. Shorthouse (2010) recognized eight ecoregions within Canada’s Prairies Ecozone—Mixed Grassland, Cypress Upland, Moist Mixed Grassland, Fescue Grassland, Tall-Grass Prairie, and Aspen Parkland. Aspen Parkland occurs at the northern limits of the grassland biome in Manitoba, Saskatchewan, and Alberta and adjacent portions of Minnesota, North Dakota, and Montana. So named because of the park-like appearance of open grassland and scattered groves of quaking aspen (*Populus tremuloides* Michx.), this zone forms an interface between the grassland biome to the south and the boreal forest to the north (Archibold 1999; Shorthouse 2010). Looman and Best (1987) recognized eastern, central, and western sections of the Parklands in the Canadian Prairie Provinces, with the herbaceous component derived from tallgrass prairie in the eastern section and mixed-grass prairie in the central and western portions.

Western Extent

While the Rocky Mountain cordillera provides an easily recognizable western physiographic boundary for the Great Plains, many species typical of Plains grassland systems extend up onto the eastern slope of the mountains, either in piedmont or foothill grassland communities or as components of the herbaceous layer of conifer woodland/savanna (Moir 1969; Mueggler & Stewart 1980; Dick-Peddie 1993; Knight 1994). For the purposes of this study, the western boundary of the study region is set at the lower elevational limits of *Pinus ponderosa* Douglas × *P. Laws.* & *C. Laws.* woodland. This delineation incorporates into the study region conifer savanna/open woodland formations that occur in places along the eastern flank of the Rocky Mountains between *Pinus ponderosa* woodland and Plains grassland systems (Knight 1999; West 1999). In the Southern Rocky Mountains, from south-central Colorado into New Mexico, this zone is occupied by juniper (*Juniperus monosperma* (Engelm.) Sarg.) or juniper-piñon pine (*Juniperus monosperma*-*Pinus edulis* Engelm.) woodland/savanna systems that also extend eastward onto the Plains in association with tablelands and canyons. Northward along the Rocky Mountain front this zone is occupied by open stands of *Juniperus scopulorum* Sarg. and/or *Pinus flexilis* E. James, which also extend eastward onto the Plains along escarpments. As with *Pinus*



FIG. 1. The Central Grassland of North America. Adapted from R. Anderson (2006). Dashed line indicates the geographical area covered in *Flora of the Great Plains* (GPFA 1986).

ponderosa savannas, the herbaceous layer of these xeric conifer woodlands is composed of grassland species, although in the northwestern part of the study region in eastern Montana the species composition has stronger affinities with Intermountain sagebrush steppe than Great Plains grasslands (Lavin & Seibert 2011).

In many places along the Rocky Mountain front, geologic strata of the Plains have been sharply upturned by the rise of the mountains into series of erosional scarps, hogbacks, mesas, and other topographic features. This rocky habitat often supports sparsely-vegetated barrens communities dominated by low-growing forbs and sub-shrubs, including many narrowly-distributed edaphic endemics. These rock outcrop communities are included in the study region.

Significant gaps in the Rocky Mountain cordillera occur in central Wyoming and central New Mexico. In both regions there exists a transition zone where the flora of the semiarid Plains gives way to species and ecological systems characteristic of arid regions to the west. In Wyoming, the transition zone is centered on the Casper Arch, an elevated area that extends from the southern tip of the Bighorn Mountains (Middle Rocky Mountains) to the northern edge of the Laramie Range (Southern Rocky Mountains). This marks the eastern edge of the Wyoming Basin, which is strongly influenced by the flora of the Great Basin (Dorn 2001). In New Mexico, the transition zone occurs between the southern end of the Sangre de Cristo Range (Southern Rocky Mountains) and the northern edge of mountains of the Basin and Range Province. This region marks a zone of tension between the flora of the Central Grassland and the flora of the Colorado Plateau (Dick-Peddie 1993). For the purposes of this study, a taxon is included in the analysis if the main part of its range lies to the east of these transition zones.

Eastern Extent

The study region encompasses the full eastern extent of the historically-known distribution of the tallgrass prairie in North America, sometimes referred to as True Prairie (Clements 1920; Rydberg 1931; Weaver 1954; Diamond & Smeins 1988) to distinguish it from mixed-grass and shortgrass prairie of the central and western Plains. Prior to almost complete conversion to agricultural production, tallgrass prairie occurred across the upper Midwest (Illinois, Iowa, Minnesota, and Wisconsin) and through the "Prairie Peninsula" (Transeau 1935) as far east as Indiana and Ohio, often in mosaic with oak woodland and savanna (Curtis 1959; Nuzzo 1986; Anderson & Bowles 1999; Will-Wolf & Stearns 1999; Cochrane & Iltis 2000; Nelson 2005).

The study region also includes the tallgrass prairies and related campestrian communities of the Ozark Highlands, which occur primarily on the Springfield Plateau of southwestern Missouri and northwestern Arkansas. Maps depicting the distribution of tallgrass prairie in North America do not usually include the Ozarks (e.g., Kuchler 1985; Lauenroth et al. 1999; Sims & Risser 2000) but the system of upland prairie, savanna, open woodland, and rock outcrop communities that historically (Bielmann & Brenner 1951; Schroeder 1982; Hanberry et al. 2014) and currently (Nelson 2005) form a continuum across the region shares a suite of grasses and other herbaceous species common to the tallgrass prairies and even mixed-grass prairies of the Central Grassland (Lawless et al. 2006). In addition, a number of species associated with rocky prairies and campestrian communities in the Ozarks also occur in association with carbonate rock outcrop habitat in Flint Hills of Kansas and Oklahoma, the Arbuckle Uplift in Oklahoma, the Blackland Prairie and Grand Prairie regions of north-central Texas, and along the eastern edge of the Edwards Plateau of central Texas. Examples include *Argyrochosma dealbata*, *Erythronium mesochoreum*, *Oenothera macrocarpa* subsp. *macrocarpa*, and *Zeltnera texensis*.

Many species characteristic of the once-extensive tallgrass prairie range beyond the study region where they occur in localized herbaceous communities within an otherwise forested landscape. On the upper Gulf Coastal Plain in Arkansas and Louisiana, these communities are remnant or naturally-isolated small-patch blackland prairies, clay prairies, saline prairies, and glades (Foti et al. 2003; MacRoberts et al. 2002, 2003, 2009, 2011; Reid et al. 2010; Foti & Witsell 2013). Farther east in Kentucky, Tennessee, Mississippi, Alabama, Georgia, and Florida, a number of species with tallgrass prairie affinities have disjunct occurrences in association with calcareous glades, barrens, and riverscours habitat (Bridges & Orzell 1986; Allison & Stevens 2001; Pringle & Witsell 2005; Lawless et al. 2006; Johnson et al. 2013; Noss 2013). Examples include *Echinacea*

pallida (Nutt.) Nutt., *Muhlenbergia cuspidata* (Torr. ex Hook.) Rydb., *Polytaenia nuttallii* DC., *Silene regia* Sims, *Silphium laciniatum* L., and *Spiranthes magnicamporum* Sheviak. Species such as these that are characteristic components of tallgrass prairie but range considerably beyond the study region are excluded from the analysis.

The grasslands and related campestran communities of the southeastern United States have recently received attention for their biological richness and critical need for conservation efforts (Noss 2013). These communities often support locally rare species, both disjunct populations of species with tallgrass prairie affinities and narrowly distributed endemics. Many of these southeastern endemics are of genera, subgeneric sections/series, or phylogenetic lineages well represented in the Central Grassland including *Astragalus bibullatus* Barneby & E.L. Bridges, *Dalea cahaba* J.R. Allison, *Pediomelum piedmontanum* J.R. Allison, M.W. Morris, & A.N. Egan, and *Physaria globosa* (Desv.) O’Kane & Al-Shehbaz. While the floristic affinities of these communities indicate past connections with the Central Grassland, the associated endemics are generally highly localized in the southeastern United States and are excluded from the present analysis.

Southern Extent

The southern reach of the study region extends considerably beyond that treated in the *Atlas of the Flora of the Great Plains* (GPFA 1977) and *Flora of the Great Plains* (GPFA 1986) to allow for consideration of the full extent of ecological systems (i.e., tallgrass, mixed-grass, and shortgrass prairie) that range from the central Plains (Kansas and Oklahoma) south into Texas. As delineated by Griffith et al. (2004) in Ecoregions of Texas, the study region includes the Texas Blackland Prairies, Cross Timbers (including the Grand Prairie and Limestone Cut Plain), Central Great Plains, Southwestern Tablelands, High Plains, and Edwards Plateau (including the Llano Uplift).

The study region excludes the Post Oak Savanna and Coastal Prairies of the Texas Coastal Plain. Recent studies have shown the Western Gulf Coastal Plain to be a distinctive floristic region and center of endemism within the larger Coastal Plain Floristic Province of North America (Estill & Cruzan 2001; Sorrie & Weakley 2001; MacRoberts et al. 2002; Noss et al. 2015). Although the Texas Blackland Prairies ecoregion is also located on the Coastal Plain, it is included in the study region because it shares many species with tallgrass prairie formations to the north (Diamond & Smeins 1985, 1988) and a number of species associated with the Edwards Plateau and Grand Prairie regions to the west have outlying occurrences in the Blackland Prairies region on outcroppings of the calcareous Austin Chalk.

On the southwest, the study region encompasses the full extent of the Llano Estacado, a vast tableland and remnant of the ancient High Plains surface that stretches from the Canadian River south to the valley of the Pecos River. Only the northern one-third of the Llano Estacado (the Texas Panhandle and adjacent part of New Mexico) was included in the region treated in the *Flora of the Great Plains* (1986) (see Fig. 1). Several unique habitats and ecological systems are associated with the Llano Estacado, notably escarpment breaks, canyonlands, playa wetlands, and dune fields (Rowell 1967; Bolen et al. 1989; Muhs & Holliday 2001; Smith 2003; Flores 2010).

MATERIALS AND METHODS

The distribution data needed to identify endemic plants in the flora of the Central Grassland were drawn from a wide variety of sources including regional, state, and local floras, floristic atlases, published descriptions of new species, recent phylogenetic literature, monographs, conservation assessments of rare species, state natural heritage program databases, examination of herbarium specimens, and the author’s 30 years of field experience in the study region. Ecological community affiliations of these taxa were drawn from the same sources.

Definition of Endemic

Following the criteria used in other phytogeographic studies (Sorrie & Weakley 2001; MacRoberts et al. 2002), a taxon is considered endemic to the study region if at least 90% of its geographic distribution is limited to the Central Grassland as here delineated. This allows for inclusion of taxa with distributions centered in the Central Grassland but extending to some extent beyond, either as a slight overlapping of its contiguous range

or as relatively few isolated, disjunct occurrences. Examples of such taxa include *Penstemon haydenii*, all but limited to the Nebraska Sandhills except for highly disjunct populations in sand habitat in the Wyoming Basin, *Castilleja genevievana*, endemic to the southwestern Edwards Plateau of Texas except for one closely adjacent occurrence in Coahuila, Mexico, and *Solidago gattingeri*, limited to rocky prairies and glades in the Ozark Plateaus of Missouri and Arkansas except for disjunct occurrences in the cedar glade region of central Tennessee.

Only taxa associated with campestrian plant communities were included in the analysis of endemism in the study region. As defined herein, "campestrian" encompasses grasslands as well as savanna-open woodland systems, shrub-steppe, and rock outcrop communities that possess a significant component of grasses, graminoids, and forbs characteristic of the regional grassland flora. This definition excludes taxa which may be geographically restricted to the study region but are ecologically associated with closed-canopy forests.

Taxonomic Considerations

The list of Central Grassland endemics presented in Appendix 1 includes a relatively small number of taxa which have been synonymized by some authorities or are of otherwise uncertain taxonomic status. These are included because they appear to be associated with a distinct region or habitat within the Central Grassland and further study may prove them worthy of recognition. Examples include *Echinocereus coccineus* subsp. *roemeri* of granite habitat in the Llano Uplift region of Texas and *Thelypodium wrightii* var. *oklahomensis* of the Mesa de Maya region of Colorado and Oklahoma. Authorities for scientific names are given in the text only for taxa not listed in Appendix 1.

Conservation Status

A taxon is here considered a plant of conservation concern if it has been given a NatureServe global conservation status ranking of G1/T1, G2/T2, or G3/T3. G-rankings pertain to species, T-rankings to subspecific taxa. Table 2 presents the NatureServe global conservation status rankings used in this study.

RESULTS AND DISCUSSION

This paper identifies 382 plant taxa that are endemic to the Central Grassland of North America as defined herein. There are 300 endemic species, eight of which have multiple subspecific taxa (totaling 10 taxa) endemic to the region (e.g., *Oenothera macrocarpa* and *Physaria ovalifolia*), and 72 endemic subspecies/varieties of more widely distributed species. Appendix 1 presents an annotated list of these taxa. Families and genera with the largest number of endemic taxa in the Central Grassland are shown in Table 3 and Table 4, respectively.

DISTRIBUTION OF ENDEMIC PLANTS IN THE CENTRAL GRASSLAND

Analysis of the geographic distribution patterns of CG endemics reveals regional concentrations of endemic taxa. These nine areas are here recognized as centers of endemism for the Central Grassland. Figure 2 shows the location of these areas within the study region and Table 5 summarizes the number of CG endemics with distributions limited to or primarily centered in each center of endemism. These taxa may range to some extent into adjacent regions but are mostly localized in a particular area of endemism and associated with distinctive habitat characteristic of the area.

The flora of these centers of endemism typically includes additional taxa that are endemic to the Central Grassland but range more widely. These are not included in Table 5 but are noted in the discussion of each center of endemism that follows. The ecological character and associated taxa of each center of endemism are presented below. Plants of conservation concern are noted by their NatureServe global conservation status ranking (see Table 2).

Arkansas Valley Barrens.—Rocky habitat along the middle Arkansas River valley in south-central Colorado between Pueblo (Pueblo County) and Cañon City (Fremont County) has long been recognized for supporting a distinctive flora with a number of rare and endemic species (Kelso et al. 1996; Colorado Native Plant Society 1997; D. Anderson 2006; Neid & Handwerk 2007; Neely et al. 2011; Weber & Wittmann 2012). This habitat has been termed the Niobrara Chalk Barrens because the regional endemics are strongly

TABLE 2. NatureServe global conservation status rankings used in this study.

G1 Critically imperiled:	At very high risk of extinction due to extreme rarity (often 5 or fewer populations), very steep declines, or other factors.
G2 Imperiled:	At high risk of extinction due to very restricted range, very few populations (often 20 or fewer populations), steep declines, or other factors.
G3 Vulnerable:	At moderate risk of extinction or extirpation due to a restricted range, relatively few populations (often 80 or fewer populations), recent and widespread declines, or other factors.
GH Possibly extinct:	Missing; known from only historical occurrences but still some hope of rediscovery.

TABLE 3. Families with the largest number of endemic taxa in the Central Grassland.

Asteraceae	79 taxa in 40 genera
Fabaceae	56 taxa in 16 genera
Brassicaceae	20 taxa in 8 genera
Onagraceae	16 taxa in 1 genus
Polygonaceae	13 taxa in 2 genera
Plantaginaceae	12 taxa in 2 genera
Polemoniaceae	11 taxa in 2 genera

TABLE 4. Genera with the largest number of endemic taxa in the Central Grassland.

<i>Astragalus</i> (Fabaceae)	18
<i>Oenothera</i> (Onagraceae)	16
<i>Physaria</i> (Brassicaceae)	13
<i>Eriogonum</i> (Polygonaceae)	12
<i>Pediomelum</i> (Fabaceae)	11
<i>Penstemon</i> (Plantaginaceae)	11
<i>Phlox</i> (Polemoniaceae)	10
<i>Dalea</i> (Fabaceae)	9
<i>Yucca</i> (Agavaceae)	7
<i>Asclepias</i> (Asclepiadaceae)	6
<i>Callirhoe</i> (Malvaceae)	6

associated with the Niobrara Formation, a Cretaceous sedimentary bedrock complex forming shale, limestone, and chalk outcrops (Scott & Cobban 1964). The most botanically important barrens occur on the Middle Chalk and Upper Chalk units of the Smoky Hill Member of the Niobrara Formation (Kelso et al. 2003).

The distributions of five taxa are limited to or primarily centered in the Arkansas Valley Barrens (Table 6). *Mentzelia chrysantha* and *Oenopsis puebloensis* are strictly limited to the Pueblo-Cañon City area. The distributions of *Mirabilis rotundifolia*, *Parthenium tetraeuris*, and *Penstemon versicolor* are concentrated in this area but also include outlying occurrences in the Raton Tablelands to the southeast where they are associated with similar calcareous rock formations (Neid et al. 2007). All of these species are of conservation concern. Three CG endemics with distributions centered on the Raton Tablelands—*Oenothera harringtonii* (G3), *Penstemon auriberbis*, and *Physaria calcicola* (G2)—also have occurrences in the Arkansas Valley Barrens. All of these species occur in sparsely vegetated (10–20% cover) rock outcrop communities dominated by low-growing perennial herbs in mosaic with sparse shrubland and piñon-juniper (*Pinus edulis* – *Juniperus monosperma*) woodland (Kelso et al. 1996; Neid & Handwerk 2007; Neely et al. 2011).

These rock outcrop communities also host a number southwestern species that are at the northern limits of their distributions in the Pueblo-Cañon City area including *Artemisia bigelovii* A. Gray, *Dalea jamesii* (Torr.) Torr. & A. Gray, *Eriogonum lachnogynum* Torr., *Frankenia jamesii* Torr. ex A. Gray, *Physaria fendleri* (A. Gray) O’Kane & Al-Shehbaz, and *Townsendia fendleri* A. Gray (G3). The distribution of *Frankenia jamesii* in the Plains is essentially bi-centric between the Arkansas Valley Barrens and the Raton Tablelands (Decker 2007; Snow 1990). This shrub occurs on calcareous rock outcrops in both regions and is the dominant species of an



FIG. 2. Centers of plant endemism in the Central Grassland. **Ark**= Arkansas Valley Barrens; **Edw** = Edwards Plateau; **LEst** = Llano Estacado Escarpments; **LUpl** = Llano Uplift; **Mesc** = Mescalero-Monahans Dunes; **Niobr** = Niobrara-Platte Tablelands; **Raton** = Raton Tablelands; **Red** = Red Bed Plains; **Rev** = Reverchon Rocklands.

TABLE 5. Centers of endemism in the Central Grassland.

Center of endemism	States of occurrence	Number of localized CG endemics
Arkansas Valley Barrens	CO	5
Edwards Plateau	TX	59
Llano Estacado Escarpments	NM, OK, TX	7
Llano Uplift	TX	11
Mescalero-Monahans Dunes	NM, TX	7
Niobrara-Platte Tablelands	CO, NE, WY	10
Raton Tablelands	CO, NM, OK	15
Red Bed Plains	KS, OK, TX	8
Reverchon Rocklands	OK, TX	11

TABLE 6. Regional endemics of the Arkansas Valley Barrens.

<i>Mentzelia chrysantha</i> G2
<i>Mirabilis rotundifolia</i> G2
<i>Oenopsis puebloensis</i> G2
<i>Parthenium tetraeuris</i> G3
<i>Penstemon versicolor</i> G3

unusual and geographically limited shrubland with which several taxa of rare and endemic plants are associated (Neid et al. 2007; Shaw et al. 1989a). Populations of *Asclepias uncialis* Greene (G3) occur within the Arkansas Valley Barrens region in association with shortgrass prairie (Decker 2006).

Edwards Plateau.—The Edwards Plateau is a vast tableland bounded on its southern and southeastern margins by the Balcones Canyonlands, a highly dissected landscape of steep canyons, narrow divides, and high-gradient streams. The Balcones Escarpment provides a sharp demarcation between the Edwards Plateau and the Gulf Coastal Plain. The central and western portion of the plateau is a broad, gently rolling plain that grades gradually from east to west into the Grand Prairie, Rolling Plains, and High Plains regions to the north. Limestones of the Edwards Formation (Cretaceous) dominate the central and western portions of the plateau. In the Balcones Canyonlands, erosion has in many places removed rocks of the Edwards Formation and exposed older Cretaceous strata, primarily the Glen Rose Formation.

Rocky habitat is a characteristic feature of the Edwards Plateau landscape. The nearly horizontal limestone strata of the Edwards Formation are relatively massive and resistant to erosion and exposures occur in the landscape as ground-level expanses of fractured bedrock and as ledges, bluffs, or rimrock above watercourses. The Glen Rose Formation is composed of alternating bands of hard limestone and soft marl and the different weathering and erosion rates of these rock types results in “stair step” topography of relatively flat-topped hills and upland divides with moderate to steeply sloping hillsides (Riskind & Diamond 1988).

Such rocky habitat plays a significant role in structuring vegetation on the Edwards Plateau. In canyons along the Balcones Escarpment, a gradation of microhabitats occurs from rimrock and cliff face to canyon bottom. Springs and seeps develop in these canyons at the contact between strata of the Edwards Formation and the lower (older) Glen Rose Formation, creating mesic conditions that support closed-canopy deciduous forest (Riskind & Diamond 1988). In contrast, xeric conditions prevail on upland sites where hard-bedding limestone strata have been exposed and soils are very shallow and rocky. Such habitat supports small patch communities of herbaceous vegetation within the surrounding matrix of juniper-oak woodland and savanna (Terletzky & Van Auken 1996; Van Auken 2000).

The Edwards Plateau is recognized as a distinct vegetational area and floristic province with one of the highest levels of vascular plant endemism of any region in North America (Correll & Johnston 1970; Poole et al. 2007). While the Edwards Plateau is included in geologic and physiographic delineations of the Great

Plains (Fenneman 1931; Hunt 1974), the floristic linkages with the Central Grassland are complicated. A high percentage of strict Edwards Plateau endemics occur in forested, relatively mesic habitat in the Balcones Canyonlands along the southern and southeastern margins of the plateau (Amos & Rowell 1988; Poole et al. 2007; Carr 2009), and analysis of phylogeographic relationships indicate strong connections with the flora of southeastern North America and the Madrean region of northern Mexico (Saghatelian 2015). Yet most of the central and western portions of the Edwards Plateau, as well as upland divides in the Canyonlands region, historically supported a mosaic of grassland with juniper and/or oak woodland or open woodland and savanna with the herbaceous layer composed of species characteristic of adjacent or interdigitating Plains grassland formations, grading from shortgrass prairie on the west to mixed-grass prairie on the east (Fowler & Dunlap 1986).

Survey of floristic, ecological, and rare plant literature for this study identified 82 taxa with distributions limited to or primarily centered on the Edwards Plateau. Of these, 21 are associated with mesophytic forest in the Balcones Canyonlands where the tree canopy is generally closed and the herbaceous layer lacks grassland species. These canyon species, including *Carex edwardsiana* E.L. Bridges & Orzell, *Clematis texensis* Buckley, *Galactia watsoniana* W.C. Holmes & Singhurst, *Lithospermum helleri* (Small) J. Cohen, *Prenanthes carrii* Singhurst, O'Kennon, & W.C. Holmes, and *Salvia pentstemonoides* Kunth & Bouché, are excluded from the present analysis. Also excluded are *Donrichardia macroneuron* (Grout) H.A. Crum & L.E. Anderson, an aquatic moss, and *Zizania texana* Hitchc., an aquatic grass, both endemic to spring-fed streams at the edge of the Edwards Plateau.

Exclusion of these forest and aquatic species leaves 59 taxa that are regional endemics of Edwards Plateau campestrian plant communities (Table 7). An additional 32 CG endemics also occur in campestrian plant communities on the Edwards Plateau but range more widely, most often into the Grand Prairie region of north-central Texas (see discussion of Reverchon Rocklands that follows).

Quite a few Edwards Plateau endemics have limited distributions and are of conservation concern, including almost all of those associated mesophytic forests in the Balcones Canyonlands. Another concentration of rare species occurs in xeric rock outcrop habitat in the semi-arid southwestern Edwards Plateau, including *Dalea sabinalis* (possibly extinct), *Eriogonum nealleyi* (G2), *Forsellesia texensis* (G1) *Perityle warnockii* (G1), and *Pomaria brachycarpa* (G2).

Llano Estacado Escarpments.—The Llano Estacado is the portion of the High Plains south of the Canadian River in northwestern Texas and eastern New Mexico. This phenomenally flat tableland is an alluvial plain composed of deposits carried eastward by ancient streams flowing from the Rocky Mountains. The surface geology of the region is shaped by underlying deposits of the Ogallala Group (Pliocene), known colloquially as “caprock” due to the tendency of its constituent sand and gravel deposits to be cemented by calcium carbonate into erosion-resistant ledges or “mortar beds” (Reeves 1972).

Much of the plateau is bounded by dramatic escarpments—the Caprock Escarpment on the east, the Mescalero Ridge on the west, and the Canadian Breaks on the north (Flores 2010). Along the length of these escarpments, erosion by headward-cutting streams has resulted in a zone of rugged, broken terrain that includes cliffs, breaks, badlands, mesas, and canyons. Where streams have carved most deeply into the surface of the plateau, as in the spectacular Palo Duro Canyon near Amarillo, Texas, older rocks of Jurassic, Triassic, and Permian systems are exposed below the Ogallala caprock, some of which have formations containing dolomite and gypsum (Hood & Underwood 2001; Nesom & O'Kennon 2008; Spearing 1991).

This center of endemism roughly corresponds to the Llano Estacado Caprock Escarpments and Breaks Shrubland and Steppe ecological system of NatureServe but encompasses escarpment habitat along the entire extent of the Llano Estacado, not just the eastern Caprock. The matrix vegetation of this rugged habitat is sparse shrubland and juniper woodland in mosaic with grassland (Rowell 1967).

The distributions of seven taxa are limited to or primarily centered in the Llano Estacado Escarpments region (Table 8). Most of these occur along the eastern Caprock Escarpment and all are associated with rock outcrop habitat. A number of other CG endemics also occur in this habitat in the region, including *Astragalus*

TABLE 7. Regional endemics of Edwards Plateau campestrian plant communities.

<i>Agalinis edwardsiana</i>	<i>Muhlenbergia involuta</i>
<i>Amorpha roemerana</i> G3	<i>Nolina lindheimeriana</i>
<i>Argemone aurantiaca</i>	<i>Paronychia lindheimeri</i>
<i>Argythamnia aphoroides</i> G2	<i>Parthenocissus heptaphylla</i>
<i>Argythamnia simulans</i>	<i>Pectis angustifolia</i> var. <i>fastigiata</i>
<i>Astragalus crassicaulus</i> var. <i>berlanderii</i> T3	<i>Pediomelum latestipulatum</i> var. <i>appressum</i>
<i>Astragalus mollissimus</i> var. <i>coryi</i> T3	<i>Penstemon guadalupensis</i> G3
<i>Berberis swaseyi</i> G3	<i>Penstemon helleri</i>
<i>Brickellia cylindracea</i>	<i>Penstemon triflorus</i> G3
<i>Brickellia dentata</i> G3	<i>Perityle lindheimeri</i> var. <i>halimifolia</i> T3
<i>Brickellia eupatorioides</i> var. <i>gracillima</i> T3	<i>Perityle lindheimeri</i> var. <i>lindheimeri</i> T3
<i>Cardamine carrii</i>	<i>Perityle warnockii</i> G1
<i>Castilleja genevievana</i> G3	<i>Phlox villosissima</i> subsp. <i>villosissima</i>
<i>Castilleja lindheimeri</i> T3	<i>Physaria sessilis</i>
<i>Chamaesyce angusta</i>	<i>Pomaria brachycarpa</i> G2
<i>Chamaesaracha edwardsiana</i>	<i>Prunus minutiflora</i>
<i>Chaetopappa bellidifolia</i>	<i>Ruellia metziae</i>
<i>Condalia hookeri</i> var. <i>edwardsiana</i> T1	<i>Sclerocactus brevihamatus</i> subsp. <i>tobuschii</i> T3
<i>Crataegus turnerorum</i> G3	<i>Selenia jonesii</i> G3
<i>Dalea sabinalis</i> GH	<i>Seymeria texana</i> G3
<i>Daucosma laciniatum</i>	<i>Streptanthus petiolaris</i>
<i>Echinocereus milleri</i> G1	<i>Tinantia anomala</i>
<i>Elymus texensis</i> G1	<i>Triodanus coloradensis</i>
<i>Eriogonum nealleyi</i> G2	<i>Valerianella stenocarpa</i> G3
<i>Eucnide bartonioides</i> var. <i>edwardsiana</i>	<i>Verbesina lindheimeri</i>
<i>Forsellesia texensis</i> G1	<i>Vernonia guadalupensis</i>
<i>Galactia texana</i>	<i>Vitis monticola</i>
<i>Hesperaloe parviflora</i> G3	<i>Yucca reverchonii</i> G3
<i>Ipomoea costellata</i> var. <i>edwardsiana</i> T2	<i>Yucca rupicola</i>
<i>Lythrum ovalifolium</i> G3	

TABLE 8. Regional endemics of the Llano Estacado Escarpments.

<i>Eriogonum alatum</i> var. <i>glabriusculum</i>
<i>Eriogonum correllii</i> G2
<i>Forsellesia planiterium</i>
<i>Hedyotis nigricans</i> var. <i>papillacea</i> T3
<i>Oenothera macrocarpa</i> subsp. <i>incana</i> T3
<i>Solidago altiplanites</i>
<i>Townsendia texensis</i>

punicus var. *punicus* (T3), *Dalea tenuifolia*, *Oenothera coryi* (G3), *Pediomelum linearifolium*, *Physaria ovalifolia* subsp. *ovalifolia*, *Scutellaria resinosa*, and *Symphotrichum fendleri*.

Llano Uplift.—The Llano Uplift of central Texas, also known as the Central Texas Uplift and Central Mineral Region, is a region of exposed Precambrian granitic bedrock mostly surrounded by the Cretaceous limestone formations of the Edwards Plateau and Grand Prairie. Fenneman (1931) included it in the Great Plains physiographic province and it is often treated as a unique subregion or vegetational area of the Edwards Plateau (Correll & Johnston 1970; Griffith et al. 2004; Poole et al. 2007). It is here treated separately from the Edwards Plateau since the regional endemics of the Llano Uplift are strictly associated with granite and do not occur on the limestone formations in the surrounding regions.

The matrix vegetation of the Llano Uplift is oak woodland and savanna in mosaic with grassland but granite exposures are sparsely vegetated and nearly devoid of woody species (Whitehouse 1933; Walters & Wyatt 1982; Singhurst et al. 2007; Taylor & O'Kennon 2016). The 11 regional endemics of the Llano Uplift (Table 9) are all associated with granite outcrops or granitic soils. All are strictly limited to the Llano Uplift

TABLE 9. Regional endemics of the Llano Uplift.

<i>Brazoria enquistii</i> G2
<i>Campanula reverchonii</i> G2
<i>Croptilon hookerianum</i> var. <i>graniticum</i>
<i>Echinocereus coccineus</i> subsp. <i>roemeri</i>
<i>Eriogonum tenellum</i> var. <i>ramosissimum</i> T3
<i>Indigofera miniata</i> var. <i>texana</i>
<i>Isoetes lithophila</i> G2
<i>Monarda stanfieldii</i> G3
<i>Packera texensis</i> G2
<i>Tradescantia pedicellata</i> G2
<i>Valerianella texana</i> G2

textcept *Packera texensis* which has a few outlying populations to the north. While not entirely limited to Llano Uplift, the distribution of CG endemic *Phlox drummondii* var. *mcallisteri* is centered on the region but also extends into adjacent areas where sandstone bedrock has weathered into coarse arenaceous soils similar in texture to granitic sands (Locklear 2011).

The lithology and inherent weathering patterns of granite bedrock present a variety of unique microhabitats to which the regional endemics of the Llano Uplift are adapted (Whitehouse 1933; Smith 1981; Walters & Wyatt 1982; Trock & O'Kennon 2003; M. Turner 2003; Poole et al. 2007; Carr 2009). *Isoetes lithophila* is restricted to shallow ephemeral pools formed atop barren bedrock. *Eriogonum tenellum* var. *ramosissimum* and *Tradescantia pedicellata* occur in bedrock fractures. *Campanula reverchonii*, *Croptilon hookerianum* var. *graniticum*, *Packera texensis*, and *Valerianella texana* occur in shallow accumulations of gravel over bedrock. *Brazoria enquistii* and *Monarda stanfieldii* are associated sandbar deposits along streams. *Brazoria enquistii*, *Campanula reverchonii*, *Croptilon hookerianum* var. *graniticum*, and *Valerianella texana* are annuals.

The Llano Uplift shares floristic connections with the Arbuckle Uplift and Wichita Mountains of Oklahoma, two regions where granitic bedrock also is exposed. Among these are remarkably disjunct occurrences of granite specialists of the Piedmont region of the southeastern United States: *Isoetes piedmontana* (N.E. Pfeiffer) C.F. Reed has been discovered in the Llano Uplift and *Cyperus granitophilus* McVaugh (G3) in the Llano Uplift, Arbuckle Mountains, and Wichita Mountains (O'Kennon & Taylor 2015; Taylor & O'Kennon 2016). These granitic areas also host disjunct and/or peripheral occurrences of xeric-adapted fern species associated with igneous rock outcrop habitat in the southwestern United States and Mexico, including *Astrolepis integerrima* (Hook.) Benham & Windham, *Cheilanthes horridula* Maxon, *Cheilanthes lindheimeri* Hook., *Cheilanthes wootonii* Maxon, *Notholaena standleyi* Maxon, and *Pellaea wrightiana* Hook. (Smith 2010; Diggs & Lipscomb 2014).

Mescalero-Monahans Dunes.—The Mescalero-Monahans Dunes are the southernmost dune fields of the Great Plains (Muhs & Holliday 1995). This extensive belt of sand dunes is located on the east side of the Pecos River valley adjacent to the western escarpment of the southern High Plains (the Llano Estacado), the portion in New Mexico called the Mescalero dunes and the portion in Texas the Monahans dunes (Holliday 2001; Muhs & Holliday 2001). Sometimes included in floristic and ecological treatments of the arid Trans-Pecos region of Texas, these dune fields are included in the study region because the matrix vegetation is sand shinnery, a type of shrub-steppe dominated by Havard shin oak (*Quercus havardii*) that is unique to the Great Plains, ranging from the Mescalero-Monahans Dunes region north through the Llano Estacado into Oklahoma (Peterson & Boyd 1998; Dhillon & Mills 1999).

The distributions of seven taxa are limited to or primarily centered in the Mescalero-Monahans Dunes region (Table 10). All but one are associated with deep, dry, loose sand, the exception being *Cyperus onerosus* which occurs in moist to wet sand in depressions among active or partially stabilized dunes (Poole et al. 2007). Two of these species, *Panicum havardii* and *Proboscidea sabulosa*, range just across the international border into the Samalayuca Dunes of northeastern Chihuahua, Mexico (Carr 2009). Ten additional CG endemics occur in

TABLE 10. Regional endemics of the Mescalero-Monahans Dunes.

<i>Cyperus onerosus</i> G2
<i>Ephedra coryi</i> G3
<i>Eurytaenia hinckleyi</i> G3
<i>Helianthus neglectus</i> G2
<i>Mentzelia strictissima</i>
<i>Panicum havardii</i>
<i>Proboscidea sabulosa</i> G3

the regional flora, all associated with sand habitat: *Chamaesyce carunculata* (G3), *Chenopodium cycloides* (G3), *Dalea lanata*, *Dimorphocarpa candicans*, *Mimosa rupertiana*, *Oenothera cinerea* subsp. *cinerea*, *Oenothera engelmannii*, *Polansia jamesii*, *Penstemon buckleyi*, and *Yucca campestris*. The recently-described CG endemic *Evolvulus arenarius* (Harms 2014) also has occurrences in the Mescalero-Monahans Dunes region and may be a species of conservation concern.

Along with this concentration of endemic plants, the biodiverse Mescalero-Monahans Dunes region hosts the endemic dune sagebrush lizard (*Sceloporus arenicolus*) plus endemic whipscorpions, scorpions, and nine endemic insect species (Longing et al. 2014).

Niobrara-Platte Tablelands.—The Niobrara-Platte Tablelands is a center of endemism located at the northern end of the High Plains section of the Great Plains physiographic province. The region extends from the Niobrara River south to the South Platte River and encompasses western Nebraska, southeastern Wyoming, and extreme northeastern Colorado.

Strata of the Arikaree Group (Late Oligocene to Early Miocene) underly the northern portion of the region, from the Niobrara River south to the North Platte River (Diffendal 1991; Swinehart et al. 1985). The component rocks are primarily fine-grained sandstones and sandy siltstones. The ancient tableland formed by these deposits has been entrenched by the Niobrara and North Platte and their tributaries into dissected uplands, forming escarpments, bluffs, buttes, and other rough, broken terrain. In this region, the Arikaree caps escarpments like the Pine Ridge and Wildcat Hills of Nebraska and the Hat Creek Breaks of Wyoming along with dramatic erosional outliers like Scotts Bluff and Chimney Rock in Nebraska.

Younger rocks of the Ogallala Group (Miocene to earliest Pliocene) replace those of the Arikaree in the southern part of the region, which encompasses the Ogallala-capped interfluvium between the valleys of the North Platte and South Platte rivers known as the Cheyenne Tableland (McMillan et al. 2002; Maher et al. 2003). The Cheyenne Tableland extends westward to the foot of the Laramie Range of the Rocky Mountains and eastward to the confluence of the North and South Platte rivers. Significant areas of exposed rock and broken topography occur along the northern and southern escarpments of the tableland. The southern escarpment marks the boundary between the High Plains and the lower-elevation Colorado Piedmont and forms an irregular line of Ogallala-capped cliffs and erosional outliers in the Pawnee Buttes region of northeastern Colorado.

Mixed-grass prairie is the matrix vegetation of the Niobrara-Platte Tablelands (Rolfmeier & Steinauer 2010). The regional endemics mostly occur in sparsely-vegetated rock outcrop communities on ridgecrests and escarpments where bedrock is at or near the surface. Soils are very shallow sandy loams or silty loams and the sites are often subjected to strong, prolonged winds. These herbaceous communities are dominated by perennial forbs, most of which have a low, compact growth habit, with grasses contributing little to the vegetation cover (Hardy 1991; Hazlett 1998; Heidel & Handley 2004; Decker 2007; Rolfmeier & Steinauer 2010; Locklear 2014).

The distributions of 10 taxa are limited to or primarily centered in the Niobrara-Platte Tablelands (Table 11). Of these, only *Eriogonum nebraskense* and *Eriogonum pauciflorum* var. *gnaphaloides* are restricted to the region. Most have distributions that are centered in the Niobrara-Platte Tablelands with scattered occurrences to the north, south, or west where they occur in similar rock outcrop habitat. Other CG endemics occurring in

TABLE 11. Regional endemics of the Niobrara-Platte Tablelands.

<i>Astragalus hyalinus</i>
<i>Astragalus sericoleucus</i>
<i>Cuscuta plattensis</i> G1
<i>Eriogonum nebraskense</i>
<i>Eriogonum pauciflorum</i> var. <i>gnaphaloides</i>
<i>Musineon tenuifolium</i>
<i>Oenothera coloradensis</i> G2
<i>Oreocarya cana</i>
<i>Paronychia depressa</i>
<i>Parthenium alpinum</i> G3

the region are *Astragalus kentrophyta* var. *kentrophyta*, *Pediomelum linearifolium*, *Phlox andicola*, and *Physaria arenosa* var. *arenosa*.

Two of the regional endemics of the Niobrara-Platte Tablelands are not associated with rock outcrop habitat. *Oenothera coloradensis* (formerly *Gaura neomexicana* subsp. *coloradensis*) is a biennial or short-lived perennial associated with subirrigated alluvial soils on the level or slightly sloping floodplains and drainage bottoms of smaller plains streams (Fertig 2000). *Cuscuta plattensis* is a twining parasitic herb reported to attach to *Ambrosia*, *Grindelia*, *Helianthus*, *Psoralea*, and *Solidago* (Nelson 1899; Handley & Fertig 2002). This critically imperiled species is documented from a very small number of collections from southeastern Wyoming within the Niobrara-Platte Tablelands and may also occur in Nebraska and Washington (Costea et al. 2006). While not entirely limited to the Central Grassland, *Dalea cylindriceps* Barneby (G3) has scattered occurrences in the Niobrara-Platte Tablelands in sandy loam soils derived from sandstone outcrops.

A number of species that are more common to the west have peripheral or disjunct occurrences in Niobrara-Platte Tablelands rock outcrop communities including *Astragalus shortianus* Nutt. and *Scutellaria brittonii* Porter of the Rocky Mountains and *Linanthus cespitosus* (Nutt.). J.M. Porter & L.A. Johnson, *Oxytropis multiceps* Nutt., and *Phlox muscoides* Nutt. of the Wyoming Basin. Another group of Wyoming Basin species occur where fine-textured siltstones and claystones of the older White River Group are exposed in deeply dissected terrain below Arikaree or Ogallala caprock. These include *Eriogonum brevicaulis* Nutt., *Lomatium nuttallii* (A. Gray) J.F. Macbr. (G3), *Phacelia hastata* Douglas ex Lehm., and *Stephanomeria runcinata* Nutt.

Raton Tablelands.—The Raton Tablelands is a center of endemism associated with the physiographic region delineated by Fenneman (1931) as the Raton section of the Great Plains. The Raton section is an uplifted area located in southeastern Colorado, northeastern New Mexico, and the adjacent portion of the Oklahoma panhandle drained by the Arkansas River on the north and the Canadian River on the south.

The Raton section has a complex geological history that includes volcanic activity and has the greatest diversity of exposed rock formations of any part of the study area. Topographic features include high mesas, extensive dissected plateaus, deep canyons, and extinct volcanoes (Lee 1902, 1903, 1912; Fenneman 1931; Flores 2010). Among the most prominent of these landforms is the Mesa de Maya, a lava-capped plateau extending about 70 km along the Colorado-New Mexico border to its terminus in Oklahoma (where it is known as the Black Mesa). Other significant geologic features are the canyons of the Purgatoire River in Colorado and the Canadian River in New Mexico, the latter being the biggest and deepest canyon in the Great Plains.

The surface geology of the Raton section is strongly influenced by Cretaceous marine strata. Sandstone of the Dakota Formation is prevalent throughout most of the region, but the endemics are primarily associated with younger carbonate Cretaceous strata, notably Niobrara Formation chalk, Greenhorn Limestone, and various shale formations. Older rocks of Jurassic, Triassic, and Permian systems are exposed within the canyons of the Purgatoire, Dry Cimarron, and Canadian rivers where these streams have cut down below the Dakota Sandstone.

This geologically complex region supports a rich flora and a diversity of plant communities. The flora and rare plants of the region has been the subject of numerous studies (Rogers 1953, 1954; Shaw et al. 1989a, 1989b;

Naumann 1991; Clark 1996; Ladyman 2005; Neid et al. 2007; Schiebout et al. 2008; Buthod & Hoagland 2015) including floristic inventories of national grasslands in the region—Comanche National Grassland in Colorado (Hazlett 2004; Kuhn et al. 2011) and Kiowa/Rita Blanca National Grassland in New Mexico (Hazlett et al. 2009). The vegetation of the region is a complex mosaic of grassland, shrubland, and conifer woodland. Shaw et al. (1989a) recognized 26 distinct plant communities from the U.S. Army Piñon Canyon Maneuver Site in southeastern Colorado, including those associated with the slopes of canyons and with outcroppings of limestone and shale.

The distributions of 15 taxa are limited to or primarily centered in the Raton Tablelands (Table 12). All of these are associated with rock outcrop habitat, although the nature of the habitat varies. *Astragalus wittmannii*, *Frasera coloradensis*, and *Packera spellenbergii* are primarily associated with ground-level exposures of Greenhorn Limestone. *Eurybia horrida*, *Nolina greenei*, *Thelypodium wrightii* subsp. *oklahomensis*, *Solidago capulinensis*, and *Yucca neomexicana* are associated with the rocky slopes of canyons, mesas, and volcanic cinder cones. *Eriogonum aliquidum*, *Oenothera harringtonii*, and *Oonopsis monocephala* occur in association with low shale hills and the saline clay flats and alluvial fans that develop downslope from eroding shale exposures.

Mirabilis rotundifolia (G2), *Parthenium tetraeuris* (G3), and *Penstemon versicolor* (G3) are CG endemics of conservation concern that occur primarily in the Arkansas Valley Barrens of south-central Colorado but also have occurrences in the Raton Tablelands (Neid et al. 2007). A number of more widely distributed CG endemics have occurrences in rock outcrop habitat in the Raton Tablelands, including *Argythamnia humilis* var. *leiosperma*, *Forsellesia planitierum*, *Grindellia inornata* (G2), *Oonopsis engelmannii* (G3), and *Oonopsis foliosa* (G3). While not endemic to the Central Grassland, *Asclepias uncialis* (G3) historically ranged through much of the shortgrass prairie of the western Great Plains and is now limited to scattered occurrences, the largest and most viable of which occur in the Raton Tablelands (Decker 2006; Neid et al. 2007).

In his analysis of the distribution of mammals in Colorado, Armstrong (1972) recognized a Raton Faunal District which he described as “the most distinctive” region of mammal diversity in the plains of eastern Colorado. This zoogeographic area strongly corresponds to the Colorado portion of the Raton Tablelands described herein, further marking the region as a biodiversity hotspot for the Great Plains.

Red Bed Plains.—The Red Bed Plains stretch from north-central and northwestern Texas through central and western Oklahoma into south-central Kansas. Within this region, brick-red shales, siltstones, and sandstones of Permian age have been exposed by erosion. The eastern part of Red Bed Plains is characterized by gently rolling hills and broad plains, but more broken topography prevails to the west where formations containing beds of gypsum and dolomite are exposed. Escarpments, mesas, buttes, and badlands are typical of areas where gypsum and dolomite cap more easily erodible shale and sandstone formations, notably the Red Hills of Kansas (Buchanan 1984) and the Cimarron Gypsum Hills, Mangum Gypsum Hills, and Weatherford Gypsum Hills of Oklahoma (Curtis et al. 2008).

Mixed-grass prairie is the matrix vegetation of most of the Red Bed Plains (Hoagland 2000; Lauer et al. 1999). The distinctive aspects of the flora, including the regional endemics, typically occur in sparsely vegetated communities associated with gypsum-bearing rock formations (Buckallew & Caddell 2003; Hoagland & Buthod 2005; Barber 2008; Nesom & O’Kennon 2008; Caddell & Rice 2012). Gypsum is a mineral rock formed by calcium sulfate dehydrate ($\text{CaSO}_4 \cdot 2\text{H}_2\text{O}$). The mineral gypsum has a pronounced effect on plant growth (Kruckeberg 2002; Escudero et al. 2014) and, generally, the higher the gypsum content of the substrate and soil the more sparse the vegetation and less diverse the flora, although the flora often includes specialized gypsum endemics.

The distributions of eight taxa are limited to or primarily centered in the Red Bed Plains (Table 13). None of the regional endemics are obligate gypsophiles. *Oenothera macrocarpa* subsp. *oklahomensis* is strongly associated with gypsum but also occurs on other substrates. A number of more wide ranging CG endemics are frequently associated with gypsum and dolomite substrates in the region, including *Astragalus lindheimeri*, *Castilleja citrina*, *Cymopterus macrorhizus*, and *Phlox oklahomensis* (G3). The flora of the Red Bed Plains region also is enriched by species characteristic of gypsum habitats in the northern Chihuahuan Desert, including

TABLE 12. Regional endemics of the Raton Tablelands.

<i>Astragalus puniceus</i> var. <i>puniceus</i> T3
<i>Astragalus wittmannii</i> G3
<i>Eriogonum aliquantum</i> G3
<i>Eurybia horrida</i> G2
<i>Frasera coloradensis</i> G3
<i>Grindelia fastigiata</i> var. <i>revoluta</i>
<i>Nolina greenei</i> G2
<i>Oenothera harringtonii</i> G3
<i>Oenopsis monocephala</i> G2
<i>Packera spellenbergii</i> G2
<i>Penstemon auriberbis</i>
<i>Physaria calcicola</i> G2
<i>Solidago capulinensis</i> G1
<i>Thelypodium wrightii</i> subsp. <i>oklahomensis</i>
<i>Yucca neomexicana</i>

TABLE 13. Regional endemics of the Red Bed Plains.

<i>Aphanostephus pilosus</i>
<i>Callirhoe scabriuscula</i> G2
<i>Chamaesaracha darcy</i>
<i>Oenothera coryi</i> G3
<i>Oenothera macrocarpa</i> subsp. <i>oklahomensis</i> T3
<i>Oenothera triangulata</i> G3
<i>Phlox drummondii</i> subsp. <i>johnstonii</i> T3
<i>Solidago mollis</i> var. <i>angustata</i> T3

Haploesthes greggii A. Gray, *Nama stevensii* C.L. Hitchc., *Oenothera hartwegii* Benth. subsp. *filifolia* (Eastw.) W.L. Wagner & Hoch, *Phacelia integrifolia* Torr., *Psilostrophe villosa* Rydb. ex Britton, and the recently described *Oenothera gayleana* B.L. Turner & M.J. Moore (Turner & Moore 2014). *Eriogonum correllii* (G2) and *Solidago altiplantites*, two species primarily associated with the Llano Estacado Escarpments region, also have occurrences in the Red Bed Plains.

Two highly-localized regional endemics of the Red Bed Plains are associated with sandy habitat. *Callirhoe scabriuscula* occurs on deep, shifting sands of ancient and contemporary terraces of the Colorado River in Coke, Mitchell, and Runnels counties in west-central Texas (Dorr 1990; Poole et al. 2009). It is listed as an endangered species by the USFWS. *Phlox drummondii* subsp. *johnstonii* occurs in similar habitat in the watersheds of the Salt Fork and Double Mountain Fork of the Brazos River in Fisher, Kent, and Stonewall counties in the southern Texas Panhandle (Locklear 2011).

While not strictly limited to the Red Bed Plains, most known occurrences of *Aschisma kansanum* are in the Red Hills region of south-central Kansas. This fascinating moss, one of the rarest in North America, is associated with Pleistocene gravel deposits where it grows at the base of relatively clear quartz pebbles that are partially imbedded in sandy soil, the translucent pebbles appear to provide protective cover and a thin film of moisture in the dry, open habitat while also letting in enough light for plant growth (Cridland 1959; Smith 1966).

Reverchon Rocklands.—The Grand Prairie of north-central Texas (sensu Diggs et al. 1999) is a vegetational region composed of the Fort Worth Prairie to the north of the Brazos River and the Lampasas Cut Plain (Limestone Cut Plain) to the south, and is contained within the Cross Timbers ecoregion of Texas (Griffith et al. 2004). Physiographically, the Grand Prairie roughly corresponds to the Comanche Plateau, a region related geologically to the Edwards Plateau to the south but from which the limestones of the Edwards Formation have in large part been eroded away, exposing older Cretaceous formations in a dissected landscape of scarps, cuestas, and mesa-like divides (Hill 1901; Fenneman 1931; Stephens & Holmes 1989). The most significant of these

strata from the standpoint of vegetation structure and endemism are the Walnut Formation and Glen Rose Formation. These are composed of interbedded erosion-resistant limestones and less consolidated strata variously termed marl, clay, or shale. The contrasting lithification of these strata create a diversity of habitats including limestone barrens, glades, and seeps that supports small patch communities of herbaceous vegetation within the surrounding matrix of juniper-oak woodland/savanna and grassland.

The distinctiveness of the flora and vegetation of the Grand Prairie has long been recognized (Hill 1901; Dyksterhuis 1946; Diamond & Smeins 1985) and recent studies of plants associated with calcareous rock outcrop habitat in the Grand Prairie have shown the region to be a center of floristic diversity with several unique and locally-restricted plant communities (Nesom & O'Kennon 2001; Swadek & Burgess 2012; Taylor & O'Kennon 2013, 2014). A number of taxa associated with barrens and glades in the Grand Prairie region also occur in calcareous rock outcrop habitat in the Arbuckle Uplift of south-central Oklahoma (Hoagland & Johnson 2001; Nesom & O'Kennon 2001), warranting inclusion of the Arbuckles into this center of endemism.

Since the region is more expansive than just the Grand Prairie of Texas, and to avoid confusion with the Grand Prairie of Arkansas, this center of endemism is here named the Reverchon Rocklands in honor of Julien Reverchon (1837–1905), French-American immigrant to Dallas and the first botanist to collect extensively in north-central Texas (Diggs et al. 1999). Reverchon collected the types of several species that are endemic to or strongly associated with the region, notably *Dalea reverchonii*, *Muhlenbergia reverchonii*, and *Pediomelum reverchonii*.

As here delineated, the Reverchon Rocklands also encompasses the East and West Cross Timbers of north-central Texas, which border the Grand Prairie on both sides. These are the southernmost belts of the larger Cross Timbers vegetational region that stretches from north-central Texas through central Oklahoma into southeastern Kansas and is characterized by a mosaic of forest, woodland, prairie, and glade communities on soils derived from sandstone formations (Diggs et al. 1999; Hoagland et al. 1999).

The distributions of 11 taxa are limited to or primarily centered in the Reverchon Rocklands (Table 14). *Silphium albiflorum* and *Yucca pallida* have the largest distributions of the regional endemics, occurring throughout the Grand Prairie region and overlapping to some extent onto the Edwards Plateau. The rarest taxa of the region are *Yucca necopina* and *Dalea reverchonii*, both endemic to north-central Texas (Poole et al. 2007; Taylor & O'Kennon 2013), and *Echinacea paradoxa* var. *neglecta*, which is endemic to the Arbuckle Uplift. *Liatris aestivalis* and *Pediomelum reverchonii* have occurrences in both the Grand Prairie and the Arbuckles.

While almost all of the regional endemics are associated with limestone, *Yucca necopina* and *Mirabilis gigantea* occur in deep, coarse-textured soils weathered from sandstone formations in the Cross Timbers area, often in association with sandy stream terraces.

The flora of the Reverchon Rocklands includes other CG endemics and locally-rare species. A number are strongly associated with rock outcrop communities in the Reverchon Rocklands but range more widely, including *Amphiachyris amoena*, *Bouteloua pectinata*, *Callirhoe pedata*, *Dysodiopsis tagetoides*, *Hedeoma reverchonii*, *Marshallia caespitosa* var. *signata*, *Matelia biflora*, *Muhlenbergia reverchonii*, and *Paronychia virginica* var. *scoparia*. Also significant are CG endemics with distributions that are essentially shared between the Grand Prairie and Edwards Plateau, including *Centaurium beyrichii*, *Juncus texanus*, *Pediomelum hypogaeum* var. *scaposum*, *Salvia engelmannii*, and *Vernonia lindheimeri*. *Ipomoea shumardiana* (G2), *Matelea edwardsensis* (G3), and *Pediomelum cyphocalyx* (G3) are rare CG endemics with occurrences in the Reverchon Rocklands. The flora also includes remarkable disjunct occurrences of several species associated with limestone cedar glades in the southeastern United States: *Gratiola quartermanniae* D. Estes (G3), *Isoetes butleri* Engelm., and *PheMERanthus calcaricus* (Ware) Kiger (G3) (Swadek 2012; Swadek & Burgess 2012; Taylor et al. 2012; Taylor & O'Kennon 2014).

Four additional regions of the Central Grassland have notable concentrations of CG endemics, some of conservation concern.

Missouri Plateau Badlands.—In the unglaciated portion of the Missouri Plateau, interstream uplands

TABLE 14. Regional endemics of the Reverchon Rocklands.

<i>Dalea hallii</i> G3
<i>Dalea reverchonii</i> G2
<i>Echinacea paradoxa</i> var. <i>neglecta</i> T1
<i>Liatris aestivalis</i>
<i>Mirabilis gigantea</i>
<i>Pediomelum reverchonii</i> G3
<i>Phlox villosissima</i> subsp. <i>latisejala</i>
<i>Senecio quaylei</i> G1
<i>Silphium albidiflorum</i>
<i>Yucca necopina</i> G1
<i>Yucca pallida</i> G3

underlain by fine-grained sedimentary rock formations (primarily siltstones and claystones) have in places been dissected into badlands. Badland terrain consists of complex networks of barren gullies and ridges, the slopes often steep and with sharp edges. Weathered outcroppings of these substrates and the fine-textured soils derived from them present a root-inhibiting combination of high rates of erosion, high shrink-swell capacity, low organic matter, low infiltration rate, low fertility and high alkalinity, with selenium sometimes present. As a result, badlands are very sparsely vegetated. Yet a suite CG endemics are strongly associated with Missouri Plateau badland habitat, notably *Astragalus barrii* (G3), *Atriplex suckleyi*, *Eriogonum pauciflorum* var. *pauciflorum*, *Eriogonum visheri* (G3), *Oonopsis multicaulis*, *Physaria arenosa* var. *argillosa* (T3), and *Physaria brassicoides* (Beatty et al. 2003; Ladyman 2006a, 2006c; Rolfsmeier & Steinauer 2010; Charboneau et al. 2013). The most extensive areas of badland topography in the Plains are the White River Badlands of western South Dakota and the Little Missouri Badlands of western North Dakota, with significant areas also occurring in northwestern Nebraska, northeastern Wyoming, and eastern Montana.

Smoky Hills.—The Smoky Hills region of north-central Kansas and adjacent south-central Nebraska lies within the Plains Border section of the Great Plains (Fenneman 1931) where the original level surface of the High Plains has been dissected by eastward-flowing, headward-cutting streams, exposing older rock formations beneath. The rolling hills of the region are formed in relatively flat-lying strata (gently dipping to the west) of Cretaceous sedimentary rocks (Buchanan 1984), notably Dakota Sandstone, Greenhorn Limestone (Hattin 1975), and Niobrara Formation limestone and chalk (Hattin 1982), geologic units that are exposed more narrowly along the eastern flank of the Rocky Mountains as upturned scarps and hogbacks. The characteristic vegetation of the region is southern mixed-grass prairie (Lauver et al. 1999).

Three CG endemics of conservation concern are strictly limited to rock outcrop habitat in the Smoky Hills: *Oenothera macrocarpa* subsp. *fremontii* (T3) occurs throughout much of the region in association with limestone and chalk outcroppings; *Eriogonum helichrysoides* (G2) and *Eriogonum jamesii* var. *simplex* (T2) are limited to the western part of the Smoky Hills where they occur in association with badlands formed in chalk of the Smoky Hill Member of the Niobrara Formation. Other more widely distributed CG endemics occurring in the region include *Physaria ovalifolia* var. *ovalifolia*, *Scutellaria resinosa*, and *Symphyotrichum fendleri*.

The Great Plains distribution of *Clematis fremontii* S. Watson is essentially limited to the Smoky Hills. This distinctive species is fairly common in limestone uplands of the Smoky Hills, occurring in 15 counties in Kansas and three in Nebraska, but is absent from the nearby Flint Hills of eastern Kansas where Permian limestones outcrop extensively. Farther to the east, scattered occurrences of *C. fremontii* are found on dolomite glades in the Ozark Highlands of Missouri (Yatskievych 2013) and Arkansas (Graves 2016). Even more highly disjunct populations of *C. fremontii* have recently been discovered in rock outcrop habitat in southeastern Tennessee and northwestern Georgia (Horn & Shaw 2007).

Southwestern Ozarks.—A number of taxa of conservation concern are associated with rocky tallgrass prairie and related campestrian plant communities in the southwestern portion of the Ozark Highlands. Four of these, *Callirhoe bushii* (G3), *Delphinium treleasei* (G3), *Echinacea paradoxa* var. *paradoxa* (T2), and *Physaris*

filiformis (G3) are strongly associated with the Springfield Plateau in southwestern Missouri and northwestern Arkansas but also have occurrences beyond, mostly in the adjacent White River Hills region of the Ozarks. Mississippian-age limestone formations dominate the surface geology of the Springfield Plateau while dolomite of Ordovician age underlies the White River Hills (Foti & Bukenhofer 1998; Nelson 2005). Topography varies from level to gently rolling hills on the western and central Springfield Plateau (referred to as the Springfield Plain in Missouri) to more dissected and strongly sloping terrain on the eastern edge and in the White River Hills.

Historically, tallgrass prairie was the matrix vegetation across the rolling uplands of the Springfield Plateau (Schroeder 1982; Nelson 2005; Foti & Witsel 2013) but was more limited in extent toward the dissected eastern portion, occurring in mosaic with savanna and open woodland complexes or as discrete landscape units on uplands bounded by timbered stream valleys (Schroeder 1982). In the more densely forested portions of the Ozark Highlands, this vegetation is presently limited to glades—small patch herbaceous communities associated with rock outcrop habitat (Nelson & Ladd 1980; Yatskievych 1999; Baskin & Baskin 2000; Ware 2002; Nelson 2005). The intergrading complex of prairies, savannas, and open woodlands on the Springfield Plateau is recognized by NatureServe as the Ozark Prairie and Woodland ecological system. In addition to the four regional endemics named above, the flora of these rocky campestrian communities is enriched by occurrences of other CG endemics of Ozarkian affinity including *Astragalus crassicaarpus* var. *trichocalyx*, *Callirhoe digitata*, *Lithospermum subsetosum*, *Nemastylis nuttallii*, *Rudbeckia missouriensis*, *Scutellaria bushii* (G3), *Solidago gattingeri* (G3), and *Valerianella ozarkana* (G3).

Texas Blackland Prairies.—The blackland prairies of Texas are floristically distinct grassland formations within the continuum of North American tallgrass prairie, combining elements of Midwestern tallgrass prairie (True Prairie) and the Gulf Coastal Prairies of Texas and Louisiana (Diamond & Smeins 1988; Diggs et al. 1999). The calcareous clay soils that support blackland prairie are derived from underlying shales, marls, and limestone bedrock. A number of CG endemics associated with limestone rock outcrop communities on the Grand Prairie and/or Edwards Plateau have outlying occurrences in the Texas Blackland Prairies ecoregion on surfacing exposures of the calcareous Austin Chalk (Cretaceous), including *Callirhoe pedata*, *Marshallia caespitosa* var. *signata*, *Physaria densiflora*, and *Salvia engelmannii* (Diggs et al. 1999). *Astragalus reflexus* (G3) and *Liatris glandulosa* (G3) are CG endemics of conservation concern that appear to be mostly restricted to outcroppings of Austin Chalk in the Texas Blackland Prairies ecoregion. The rarest plant of the ecoregion is *Hymenoxys perpygmaea* (G1), a recently-described species (Mink et al. 2012) associated with the regionally-restricted and critically imperiled (G1) *Sporobolus silveanus* Swallen – *Carex meadii* Dewey herbaceous grassland community (Diamond & Smeins 1985, 1988, 1993). Areas of deeper soils in the Texas Blackland Prairies ecoregion have been almost entirely lost to agricultural production with the result that this grassland exists today in only a few preserves, small remnants, and native hay meadows.

ECOLOGY OF ENDEMIC PLANTS OF THE CENTRAL GRASSLAND

Of the 382 CG endemics, 299 or 78% are habitat specialists, associated with rock outcrop, sand, hydric, or riparian habitats (Table 15). These habitats typically occur as discrete, localized areas and support unique plant communities within the regionally dominant matrix vegetation of grassland, shrubland, or woodland (Kelso et al. 2001). Such “habitat islands” with their distinctive and specialized floras have been termed insular ecosystems and are often important to regional biodiversity (Cartwright & Wolfe 2016).

Rock Outcrop Habitat.—Fifty-nine percent of CG endemics (225 taxa) are primarily associated with rock outcrop habitat (indicated by the abbreviation *r* in Appendix 1). Rock outcrop habitat is the prevailing ecological association of regional endemics in all but one of the nine centers of endemism recognized above, the exception being the Mescalero-Monahans Dunes. The descriptive term “rock outcrop” is here used broadly to signify rocky, shallow-soil habitat that supports plant associations that are distinct from the surrounding matrix vegetation on zonal soils and encompasses a range of habitat from massive exposures of bedrock to areas of rocky/gravelly soils. The topographic position of this habitat is upland and includes interstream divides, escarpments, buttes, mesas, canyon slopes, etc. where bedrock is exposed.

TABLE 15. Habitat affinities of Central Grassland endemics.

Habitat	Number of taxa
Rock outcrop	225
Sand	53
Hydric	14
Riparian	7
	299 Total habitat specialists

The majority of CG rock outcrop endemics are associated with limestone or other carbonate substrates. Limestone formations dominate the surface geology of the Edwards Plateau and Reverchon Rocklands, two major centers of endemism for the Central Grassland, as well as the Springfield Plateau of the southwestern Ozark Highlands. Elsewhere in the Central Grassland, more geographically-limited exposures of limestone also support CG endemics, notably outcroppings of the Greenhorn Limestone in the Raton Tablelands and Smoky Hills. A number of narrowly-distributed endemics of conservation concern are associated with chalky limestone strata of the Niobrara Formation in the Arkansas Valley Barrens region and the Smoky Hills.

The most geographically-restricted CG endemics are those associated with granite. Within the study region, granite exposures occur in the Llano Uplift of Texas, the Arbuckle Mountains and Wichita Mountains of Oklahoma, the Hartville Uplift in Wyoming, the Black Hills in South Dakota and Wyoming, and certain isolated mountain ranges on the plains of eastern Montana (Diffendal 2017). Granite specialists are typically associated with unique microhabitats that result from the lithology and inherent weathering patterns of granite bedrock and are often of conservation concern due to their limited distribution and narrow ecological association (Shure 1999). Eleven taxa are limited to granite substrates in the Llano Uplift, nine of which are of conservation concern. Two additional CG endemics, *Phlox pilosa* subsp. *longipilosa* (T2) and *Echinocereus reichenbachii* subsp. *baileyi* (T3) are associated with granite substrates in the Wichita Mountains.

Aside from granite specialists, substrate fidelity is not pronounced among CG rock outcrop endemics. Within its limited range in southeastern Colorado, *Frasera coloradensis* occurs in association with a variety of Cretaceous strata including Greenhorn Limestone, Graneros Shale, and Dakota Sandstone (Naumann 1981). *Phlox oklahomensis* occurs in association with sandstone, shale, gypsum, and dolomite of the Ogallala, Rush Springs, and Marlow formations in the Gypsum Hills of Kansas and Oklahoma and limestone of the Permian-age Chase and Council Grove groups in the southern Flint Hills of Kansas (Springer and Tyrll 2003). The narrowly-distributed *Mentzelia chrysantha* (D. Anderson 2006) and *Physaria bellii* (Colorado Natural Heritage Program 1997) of Colorado are strongly associated with Niobrara Formation chalk but also occur on other strata where these are exposed in proximity.

This general broad tolerance of different rock types or lack of fidelity to a specific geologic substrate has also been observed among rock outcrop endemics in the eastern United States (Baskin & Baskin 1988) and suggests that chemical properties of the parent material are not a significant factor in the association of CG endemics with rock outcrop habitat. This conclusion is supported by the work of Kelso et al. (2003) in which they investigated factors influencing plant endemism associated with the Smoky Hill Chalk Member of the Niobrara Formation in the Arkansas Valley Barrens of Colorado and determined that structural components of the barrens (e.g., dry, unstable slopes and extremely shallow, mineralized soils) were more influential than the chemistry of the bedrock and that the endemics were functionally adapted to survive in habitat that is stressful enough to exclude more competitive vegetation.

As geobotanist Arthur Kruckeberg (1986) observed, "Though a region's climate sets the limits for a biota, geology enriches local discontinuity and habitat diversity." The common perception of the Central Grassland as a region of subdued landscape diversity and broad environmental gradients is generally true in the eastern portion of the region, particularly where the landscape is mantled by deposits of loess and glacial till. But exposures of surfacing bedrock in the central and western portions of the Central Grassland (Fenneman 1931;

Diffendal 2017) significantly increase local landscape heterogeneity and support more structurally complex vegetation and higher overall plant diversity than the surrounding matrix vegetation (Kelso et al. 2001). University of Kansas biogeographer P.V. Wells (1970) was correct in noting that “contrary to popular opinion, the plains are by no means lacking in edaphic diversity,” but went on to conclude, “Nevertheless, the considerable potential for the evolution of narrow endemics has not been realized in the existing flora of the Central Plains region.” An additional four decades of field work has yielded a much more complete knowledge of the flora of the region and has allowed the present analysis which reveals a strong correlation between geology and endemism in the Central Grassland.

Sand Habitat.—Fifty-three CG endemics are primarily associated with sand habitat (indicated by the abbreviation *s* in Appendix 1). Eolian sand sheets and sand dunes are common landforms in the central and western Plains (Muhs & Holliday 1995). At present, these dune fields are mostly stabilized by vegetation. Some Plains dune fields are expansive, notably the Nebraska Sandhills at approximately 50,000 sq. km. Extensive dune fields also occur along the South Platte River in Colorado and in the Pecos River valley adjacent to the western escarpment of the Southern High Plains in New Mexico and Texas. Smaller, more isolated dune fields and belts occur in the Plains where loose sand has been blown into dunes from alluvial deposits along streams that have traversed areas of sandstone bedrock. This type of habitat is particularly common below the eastern escarpment of the southern High Plains in Oklahoma and Texas (Muhs & Holliday 1995, 2001; Curtis et al. 2008) where eastward-flowing streams have downcut into sandstone rocks of the Ogallala caprock and older (deeper) formations of the Jurassic, Triassic, and Permian systems. Where these areas of sand occur as discrete belts or “islands” within the surrounding landscape of non-sandy or less sandy soil types, they enrich the local flora with arenaceous species (Ramaley 1939; Kelso et al. 2007; Nesom & O’Kennon 2008).

Some CG sand endemics have relatively large areas of distribution in the western Plains, including *Astragalus ceramicus* var. *filiformis*, *Dalea arenicola*, *Eriogonum annuum*, *Ipomoea leptophylla*, *Mentzelia nuda*, *Phlox andicola*, and *Polanisia jamesii*. Others, notably *Callirhoe scabriuscula* and *Phlox drummondii* subsp. *johnstonii* of Texas, are much more limited in distribution and are of conservation concern. Most CG sand endemics are species of the southern High Plains and are associated with shrub-steppe communities, either sandsage prairie dominated by *Artemisia filifolia* Torr. or shinnery dominated by *Quercus havardii*. Examples of these include *Aphanostephus pilosus*, *Chamaesyce carunculata*, *Chenopodium cycloides*, *Euphorbia strictior*, *Oenothera cinerea* subsp. *cinerea*, *Oenothera engelmannii*, and *Penstemon buckleyi*. The greatest concentration of CG sand endemics occurs in the Mescalero-Monahans Dunes of New Mexico and Texas, the southernmost dune fields of the Great Plains where seven regional endemics occur.

Although the Nebraska Sandhills is the largest dune field in the Western Hemisphere, no vascular plants are strictly limited to this region. The critically imperiled *Penstemon haydenii* was long thought to occur only in the Nebraska Sandhills until an outlying population center was discovered in 1996 in the Ferris Dune Field of central Wyoming (Heidel 2012).

The sandy habitat that hosts localized endemics in the western Plains is primarily derived from the activity of wind (eolian processes). But in the Cross Timbers region of north-central Texas and south-central Oklahoma, CG endemics *Ipomoea shumardiana*, *Mirabilis gigantea*, *Monarda punctata* var. *intermedia*, and *Yucca necopina* are associated with deep, coarse-textured soils weathered from sandstone strata by colluvial and/or alluvial processes.

The association of distinct floristic assemblages, often including endemic taxa, with geographically isolated areas of sandy habitat is well documented. Examples in the United States include the Fall-line Sandhills of the Carolinas and Georgia (Sorrie & Weakley 2001), the Lake Wales Ridge of peninsular Florida (Menges 1999), and the Carrizo Sands of east-central Texas (Diggs et al. 2006). In a study of endemism in the flora of the West Gulf Coastal Plain of Texas, 53% of the 96 endemics were associated with xeric sandylands (MacRoberts et al. 2002).

Hydric and Riparian Habitats.—Fourteen CG endemics are primarily associated with hydric habitat (indicated by the abbreviation *h* in Appendix 1). Wetlands and hydric habitats of various kinds occur through-

out the study region, and provide discrete landscape patches that enhance heterogeneity and species diversity. The majority of vascular plants associated with hydric features in the Central Grassland are relatively widespread, obligate or facultative wetland generalists, with endemics being relatively uncommon in this habitat type.

One of the most unique landscape features of the study region are the freshwater playa wetlands of the southern and central Great Plains. Playas are shallow, closed basin, precipitation-filled recharge wetlands. The greatest number and highest concentration of playas occur in the southern High Plains on the Llano Estacado of northwestern Texas and eastern New Mexico where they are the most significant topographic feature of the region and provide islands of biodiversity in one of the most intensively agriculturalized areas of North America (Bolen et al. 1989; Hoagland & Collins 1997; Smith 2003). Nearly 350 species of vascular plants have been identified in association with playas in the Southern High Plains (Rowell 1971; Haukos & Smith 1997). Several of these are relatively wide-ranging CG endemics, including *Ambrosia grayi*, *Oenothera canescens*, and *Suckleya suckleyana*. *Polygonum texense* is endemic to Texas where it occurs in seasonally moist habitat on the southern High Plains and the adjacent northwestern Edwards Plateau.

Less well-known concentrations of playas occur in the central Great Plains, primarily in western Kansas and eastern Colorado (Cariveau & Johnson 2007; Evans 2010). These playas differ from those on the Llano Estacado in generally being smaller and shallower and experiencing shorter periods of inundation, and with vegetation dominated by graminoids tolerant of wet periods (e.g., *Elymus smithii* (Rydb.) Gould and *Buchloe dactyloides* (Nutt.) Engelm.) as opposed to wetland species. These playas are also more likely to occur in the context of relatively intact grassland communities than those of the intensively farmed Llano Estacado. *Ambrosia linearis* is endemic to east-central Colorado and is closely associated with playa habitat, occurring at the outer edge of the zone of inundation (Locklear 1989, 1990). The highly restricted distribution of *A. linearis* is remarkable in a genus known for widespread weeds.

Closed basin wetlands similar to High Plains playas occur on the western edge of the Edwards Plateau and have been designated the Edwards Plateau Playa ecological system by NatureServe. These shallow wetlands are internally draining depressions of karstic origin on level plateau surfaces, forming from solution of underlying limestone formations. *Selenia jonesii*, an annual, is endemic to this habitat.

A number of CG endemics are associated with groundwater seeps in the Reverchon Rocklands. The surface geology of this region is dominated by interbedded limestone and marl formations and seeps occur on hillslopes where groundwater flowing through fractured limestone encounters an impervious rock layer and flows laterally. Such sites experience alternating vernal saturation and summer desiccation and support herbaceous vegetation of which several Plains endemics are characteristic species (Swadek & Burgess 2012). *Muhlenbergia reverchonii* is typically a component of these communities and is often the dominant species present, hence the colloquial name “Muhly seep” for this vegetation (Dyksterhuis 1946; Llado & Slattery 2015). Other CG endemics characteristic of this habitat include *Bouteloua pectinata*, *Eleocharis occulta*, *Juncus texanus*, and *Rhynchospora nivea*. Groundwater seep habitat occurs in the context of xeric rock outcrop habitat (glades and barrens) that supports additional CG endemics. This unique vegetation also occurs in limestone areas of the Arbuckle Uplift of Oklahoma (Hoagland & Johnson 2001), which is included as part of the Reverchon Rocklands.

Oenothera coloradensis (formerly *Gaura neomexicana* subsp. *coloradensis*) is a noteworthy CG endemic associated with hydric habitat. This short-lived perennial is associated with subirrigated alluvial soils on the level or slightly sloping floodplains and drainage bottoms of smaller plains streams (Fertig 2000). The largest populations of this imperiled taxon occur in southeastern Wyoming, with its historical range extending into adjacent parts of Colorado and Nebraska.

Seven CG endemics are primarily associated with riparian habitat (indicated by the abbreviation RIP in Appendix 1), particularly gravel and cobble bars along high-gradient streams as they emerge from canyons cut into the Balcones Escarpment. These often intermittent streams typically occupy only a small part of the streambed except during periods of flooding. Scouring by periodic flash floods keep this riparian corridor

habitat relatively open and free of woody vegetation. Examples of endemics associated with Edwards Plateau riverscours habitat include *Brickellia dentata*, *Crataegus turnerorum*, *Lythrum ovalifolium*, and *Phlox villosissima* subsp. *villosissima*.

CONSERVATION STATUS OF ENDEMIC PLANTS OF THE CENTRAL GRASSLAND

Of the 382 CG endemics, 124 or 33% are of conservation concern, having been given NatureServe global conservation status rankings of critically imperiled (G1/T1; 14 taxa), imperiled (G2/T2; 32 taxa), vulnerable (G3/T3; 77 taxa), or possibly extinct (GH; 1 species). Eight of these are also listed by the U.S. Fish and Wildlife Service as threatened or endangered.

Appendix 2 presents the entire list of CG endemics of conservation concern. Three additional CG endemics may be of conservation concern but are recently-described and not yet assigned conservation status rankings: *Cardamine carrii* (Turner 2012a) and *Eucnide bartonioides* var. *edwardsiana* (Turner 2012b) of the Edwards Plateau and *Evolvulus arenarius* (Harms 2014) of the southern High Plains.

Central Grassland endemics of conservation concern are not distributed uniformly across the study region. Of the 124 at-risk taxa in the Central Grassland, 78 or 63% are primarily associated with one of the centers of endemism identified in this study (Table 16). States with highest number of at-risk CG endemics are Texas (75), Colorado (22), Oklahoma (22), New Mexico (21), and Kansas (19) (Table 17).

TABLE 16. Distribution of Central Grassland endemics of conservation concern by center of endemism.

	Total	G1/T1	G2/T2	G3/T3	GH
Edwards Plateau	30	5	4	20	1
Raton Tablelands	11	1	5	5	
Llano Uplift	8	0	6	2	
Reverchon Rocklands	7	3	1	3	
Red Bed Plains	6	0	1	5	
Arkansas Valley Barrens	5	0	3	2	
Mescalero-Monahans Dunes	5	0	2	3	
Llano Estacado Escarpments	3	0	1	2	
Niobrara-Platte Tablelands	3	1	1	1	

TABLE 17. Distribution of Central Grassland endemics of conservation concern by state/province.

	Total	G1/T1	G2/T2	G3/T3	GH
Texas	75	8	16	50	1
Colorado	22	1	10	11	
Oklahoma	22	1	4	17	
New Mexico	21	1	6	14	
Kansas	19	2	4	13	
Nebraska	10	3	1	6	
Missouri	8	0	2	6	
Arkansas	7	0	1	6	
Wyoming	7	2	1	4	
South Dakota	6	0	0	6	
Iowa	4	0	1	3	
Illinois	3	0	1	2	
Minnesota	3	0	0	3	
Montana	3	0	0	3	
Wisconsin	3	0	1	2	
Indiana	2	0	1	1	
North Dakota	2	0	0	2	
Manitoba	1	0	0	1	
Michigan	1	0	0	1	
Ohio	1	0	0	1	

The centers of endemism identified for the Central Grassland could constitute or contain sites worthy of recognition as an “Important Plant Area” in that they contain assemblages of globally rare (G1-G3) taxa as well as disjunct and peripheral occurrences of plants that may be locally rare and of conservation concern at the state level (S1-S2). The identification of Important Plant Areas is used as a conservation planning tool in a number of state rare plant conservation strategies in the United States.

CONCLUSION

Temperate grasslands are among the most modified ecosystems on Earth due to productive soils and relative ease of conversion to cropland (Henwood 2010). Globally, temperate grasslands have been identified as one of the two terrestrial biomes in which biodiversity and ecosystem services are at greatest risk because of extensive habitat conversion and limited habitat protection (Hoekstra et al. 2005). In the Central Grassland of North America, 21.5 million hectares (53 million acres) of grassland, an area the size of Kansas, was converted to cropland between 2009 and 2015, with the region losing more habitat in 2014 than the Brazilian Amazon lost to deforestation (Gage et al. 2016; World Wildlife Fund 2016). Among the many consequences of this habitat loss is the continental-scale population decline of grassland birds, one of the prominent wildlife conservation crises of the 21st century (Brennan & Kuvlesky 2005).

The Central Grassland of North America encompasses all or part of 17 states and three Canadian provinces. Developing strategies and priorities for the conservation and protection of biological diversity for such a vast region is challenging. It is hoped that the findings of this study will be useful in focusing conservation action on the habitats, ecological associations, and regions of the Central Grassland that host the highest concentrations of unique and at-risk plant species and associated biological diversity.

APPENDIX 1

ANNOTATED LIST OF PLANTS ENDEMIC TO THE CENTRAL GRASSLAND OF NORTH AMERICA

The following compilation enumerates all plant taxa endemic to the Central Grassland as defined herein. Families are arranged alphabetically within major groups. Taxa are listed alphabetically within their respective families by genus, species, and subspecific epithet. Each entry is annotated with the following information where pertinent.

Distribution: States and provinces in which a taxon occurs are listed for CG endemics that are (1) localized within a center of endemism or (2) of limited distribution within the Central Grassland (three states or less). No states/provinces of occurrence are provided for CG endemics that are more widely distributed unless ranked as a plant of conservation concern. Standard abbreviations are used for states/provinces; **MX** = Mexico; **adj** = ranges into adjacent region; **disj** = disjunct occurrences. Taxa that are regional endemics of a center of endemism are noted using the following abbreviations:

Ark	Arkansas Valley Barrens	Niobr	Niobrara-Platte Tablelands
Edw	Edwards Plateau	Raton	Raton Tablelands
LEst	Llano Estacado Escarpments	Red	Red Bed Plains
LUpl	Llano Uplift	Rev	Reverchon Rocklands
Mesc	Mescalero-Monahans Dunes		

Habitat: If a taxon is specialized to a particular type of habitat identified in the study, that habitat is indicated using the abbreviations below. No habitat type is indicated if the taxon is not a specialist of one of these habitats.

H	Hydric	RIP	Riparian
R	Rock outcrop	s	Sand

Conservation status: Taxa of conservation concern are indicated by their NatureServe global conservation status ranking. G-rankings pertain to species, T-rankings to subspecific taxa. See Table 2 for more detailed descriptions of these rankings. Taxa listed by the U.S. Fish and Wildlife Service as threatened or endangered are also indicated.

G1/T1 Critically imperiled
G2/T2 Imperiled
G3/T3 Vulnerable
GH Possibly extinct

US-LT U.S. Fish and Wildlife Service Listed
 Threatened

US-LE U.S. Fish and Wildlife Service Listed
 Endangered

Other notes: Brief notes pertaining to taxonomic issues are provided where appropriate. Important references pertaining to ecology or conservation assessment are also noted.

BRYOPHYTES

Pottiaceae

Aschisma kansanum A.L. Andrews KS R G1
Molendoa ogalalensis (G.L. Smith) R.H. Zander [*Ozobryum ogalalense* G.L. Smith] KS, NE, disj MX R G1?
Pterygoneurum subsessile (Brid.) Jur. var. *kieneri* Habeeb NE G4?TNR
 | known from single collection (Chase County, NE) but recognized in FNA (vol. 27, 2007)

PTERIDOPHYTES

Isoetaceae

Isoetes lithophila Pfeiffer TX (**LUpl**) R G2

Pteridaceae

Argyrososma dealbata (Pursh) Windham R

GYMNOSPERMS

Ephedraceae

Ephedra coryi E.L. Reed NM, TX (**Mesc**) s G3

ANGIOSPERMS: MONOCOTS

Agavaceae

Camassia angusta (Engelm. & A. Gray) Blank.
Hesperaloe parviflora J.M. Coult. TX (**Edw**), adj MX R G3
Yucca campestris McKelvey NM, TX (**Mesc**) s
Yucca glauca Nutt.
Yucca necopina Shinners TX (**Rev**) s G1
Yucca neomexicana Wootton & Standl. CO, NM, OK (**Raton**) R
Yucca pallida McKelvey TX (**Rev**) R G3
Yucca reverchonii Trel. TX (**Edw**) R G3
Yucca rupicola Scheele TX (**Edw**) R

Alliaceae

Allium canadense L. var. *fraseri* Ownbey R
Allium canadense L. var. *hyacinthoides* (Bush) Ownbey & Aase
 OK, TX R
Allium canadense L. var. *lavendulare* (Bates) Ownbey & Aase R
Allium perdulce S.V. Fraser var. *perdulce* s
Allium stellatum Fras.ex Ker. Gawl.

Commelinaceae

Tinantia anomala (Torr.) C.B. Clarke TX (**Edw**) R
Tradescantia bracteata Small
Tradescantia edwardsiana Tharp TX R
Tradescantia pedicellata Celariar TX (**LUpl**) R G2
Tradescantia gigantea Rose TX, disj LA R
Tradescantia tharpianii E.S. Anderson & Woodson R

Cyperaceae

Carex missouriensis P. Rothr. & Reznicek

Carex perdentata S.D. Jones OK, TX R
Carex shinersii P. Rothr. & Reznicek AR, KS, OK, TX H G3
Cyperus onerosus M.C. Johnst. TX (**Mesc**) H G2
Cyperus setigerus Torr. & Hook. H
Eleocharis occulta S.G. Sm. OK, TX H
Rhynchospora nivea Boeckeler OK, TX H

Iridaceae

Nemastylis nuttallii Pickering ex R.C. Foster AR, MO, OK R
Sisyrinchium campestre E.P. Bicknell

Juncaceae

Juncus brachyphyllus Wiegand H
Juncus texanus (Engelm.) Coville OK, TX H

Liliaceae

Erythronium mesochoreum Knerr

Melanthiaceae

Zigadenus nuttallii (A. Gray) S. Watson R

Nolinaceae

Nolina greeneyi S. Watson ex Trel. CO, NM, OK (**Raton**) R G2
Nolina lindheimeriana (Scheele) S. Watson TX (**Edw**) R

Orchidaceae

Platanthera praeclara Sheviak & M.L. Bowles MB; IA, KS, MN, MO,
 ND, NE, OK, SD G3/US-LT

Poaceae

Bouteloua pectinata Feath. OK, TX R
Elymus texensis J.J.N. Campb. TX (**Edw**) R G1
Muhlenbergia ammophila P.M. Peterson [*Redfieldia flexuosa* (Thurb.)
 Vasey] s
Muhlenbergia involuta Swallen TX (**Edw**) R | possibly of hybrid origin
 involving *M. lindheimerii* Hitchc. × *M. reverchonii* Vasey & Scribn.
 (= *M. × involuta*)
Muhlenbergia reverchonii Vasey & Scribn. OK, TX H
Panicum havardii Vasey NM, TX (**Mesc**), adj MX s

Themidaceae

Androstegium coeruleum (Scheele) Greene KS, OK, TX

ANGIOSPERMS: DICOTS

Acanthaceae

Ruellia drummondiana A. Gray TX R
Ruellia metziae Tharp TX (**Edw**) R

Amaranthaceae

Atriplex suckleyi (Torr.) Rydb. R
Chenopodium cycloides A. Nelson CO, KS, NE, NM, TX s G3 |
 Ladyman 2006b
Chenopodium pallescens Standl. R
Suckleya suckleyana (Torr.) Rydb. H

Apiaceae

- Cymopterus macrorhizus* Buckley OK, TX R
Cymopterus montanus Torr. & A. Gray R
Daucosma laciniatum A. Gray TX (**Edw**) R
Eryngium diffusum Torr. OK, TX s
Eryngium leavenworthii Torr. & A. Gray R
Eurytaenia hincleyi Mathias & Constance NM, TX (**Mesc**) s G3
Lomatium foeniculaceum (Nutt.) J.M. Coult. & Rose subsp.
daucifolium (Torr. & A. Gray) W.L. Theobald R
Musineon tenuifolium (Nutt. ex Torr. & A. Gray) J.M. Coult. & Rose
 CO, NE, SD, WY (**Niobr**) R

Apocynaceae

- Amsonia ciliata* Walter var. *texana* (A. Gray) J.M. Coult. OK, TX R

Asclepiadaceae

- Asclepias arenaria* Torr. s
Asclepias lanuginosa Nutt.
Asclepias meadii Torr. ex A. Gray IA, IL, IN, KS, MO, WI G2/US-LT
Asclepias pumila (A. Gray) Vail
Asclepias stenophylla A. Gray
Asclepias sullivantii Engelm. ex A. Gray
Matelea edwardsensis Correll TX G3
Matelea biflora (Raf.) Woodson OK, TX R

Asteraceae

- Ambrosia grayi* (A. Nelson) Shinners H
Ambrosia linearis (Rydb.) W.W. Payne CO H G3 | Locklear 1989, 1990
Amphiachyris amoena (Shinners) Solbrig TX R
Aphanostephus pilosus Buckley OK, TX (**Red**) s
Brickellia cylindracea A. Gray & Engelm. TX (**Edw**) R
Brickellia dentata (DC.) Sch. Bip. TX (**Edw**) RIP G3
Brickellia eupatorioides (L.) Shinners var. *gracillima* (A. Gray) B.L.
 Turner TX (**Edw**) RIP G5T3
Brickellia eupatorioides (L.) Shinners var. *texana* (Shinners) Shinners R
Chaetopappa bellidifolia (A. Gray & Engelm.) Shinners TX (**Edw**) R
Coreopsis palmata Nutt.
Coreopsis wrightii (A. Gray) H.M. Parker OK, TX R
Croptilon hookerianum (Torr. & A. Gray) House var. *graniticum* (E.B.
 Sm.) E.B. Sm. TX (**Lupl**) R
Croptilon hookerianum (Torr. & A. Gray) House var. *validum* (Rydb.)
 E.B. Sm. KS, OK, TX, disj AR
Dieteria canescens Nutt. var. *nebraskana* (B.L. Turner) D.R. Morgan
 & R.L. Hartm. NE, SD s G5T3?
Diaperia prolifera (Nutt. ex DC.) Nutt. var. *prolifera*
Dysodiopsis tagetoides (Torr. & A. Gray) Rydb. OK, TX R
Echinacea angustifolia DC.
Echinacea atrorubens (Nutt.) Nutt. KS, OK, TX G3 R
Echinacea paradoxa (Norton) Britton var. *neglecta* McGregor OK
 (**Rev**) R G2T1
Echinacea paradoxa (Norton) Britton var. *paradoxa* AR, MO R G2T2
Erigeron bellidiastrum Nutt. var. *robustus* Cronquist s
Eurybia horrida (Wooton & Standl.) G.L. Nesom CO, NM (**Raton**) R G2?
Grindelia adenodonta (Steyerm.) G.L. Nesom TX R
Grindelia fastigiata Greene var. *revoluta* (Steyerm.) Adr. Bartoli &
 Tortosa [G. *revoluata* Steyerm., G. *hirsutula* Hook. & Arn. var.
revoluta (Steyerm.) Ackerf.] CO (**Raton**) R
Grindelia inornata Greene CO, NM R G2? | Bartoli & Tortosa 2012
 place under G. *fastigiata* Greene var. *fastigiata*
Grindelia lanceolata Nutt. var. *texana* (Scheele) Shinners OK, TX,
 adj MX R
Grindelia nuda Alph. Wood var. *nuda* R
Helianthus neglectus Heiser NM, TX (**Mesc**) s G2Q | possibly geo-
 graphically restricted ecologically distinct subspecies of the
 more widespread *H. petiolaris* Nutt. (Raduski et al. 2010)
Helianthus salicifolius A. Dietr. R
Heterotheca canescens (DC.) Shinners
Heterotheca stenophylla (A. Gray) Shinners [including var.
angustifolia (Rydb.) Semple]
Heterotheca villosa (Pursh) Shinners var. *ballardii* (Rydb.) Semple
Hymenopappus flavescens A. Gray var. *flavescens* s
Hymenopappus tenuifolius Pursh
Hymenoxys perpygmaea W.C. Holmes, Singhorst & Mink TX G1 |
 Mink et al. 2012
Iva corbinii B.L. Turner TX | apparently restricted to floodplain of
 Colorado River in Travis County, TX (Turner 2009); probably of
 conservation concern
Liatrix aestivalis G.L. Nesom & O'Kennon OK, TX (**Rev**) R
Liatrix glandulosa G.L. Nesom & O'Kennon TX R G3
Liatrix lancifolia (Greene) Kittel s
Liatrix punctata (Hook.) var. *mucronata* (DC.) B.L. Turner
Liatrix squarrosa (L.) Michx. var. *glabrata* (Rydb.) Geiser
Marshallia caespitosa Nutt. ex DC. var. *signata* Beadle & F.E. Boynton
 OK, TX, disj. AR, LA, MO R
Nothocalais cuspidata (Pursh) Greene
Oenopsis engelmannii (A. Gray) Greene CO, KS R G3
Oenopsis foliosa (A. Gray) Greene CO R G3
Oenopsis monocephala A. Nelson [O. *foliosa* var. *monocephala*
 (A. Nelson) Kartesz & Gandhi] CO (**Raton**) R G4T2 | Schulz &
 Shaw 1992
Oenopsis multicaulis (Nutt.) Greene R
Oenopsis puebloensis G. Brown & Evans CO (**Ark**) R G2
Packera spellenbergii (T.M. Barkley) C. Jeffrey NM (**Raton**) R G2
Packera texensis O'Kennon & Trock TX (**Lupl**) R G2
Parthenium alpinum (Nutt.) Torr. & A. Gray CO, WY (**Niobr**), disj NM
 R G3 | Heidel & Handley 2004
Parthenium tetraeuris Barneby CO (**Ark**) R G3
Pectis angustifolia Torr. var. *fastigiata* (A. Gray) D.J. Keil. TX (**Edw**) R
Perityle lindheimeri (A. Gray) Shinners var. *halimifolia* (A. Gray) A.M.
 Powell TX (**Edw**) R G4T3Q
Perityle lindheimeri (A. Gray) Shinners var. *lindheimeri* TX (**Edw**) R
 G4T3
Perityle warnockii A.M. Powell TX (**Edw**) R G1
Pyrrhopappus grandiflorus (Nutt.) Nutt. KS, OK, TX R
Rayjacksonia annua (Rydb.) R.L. Hartm. & M.A. Lane s
Rudbeckia missouriensis Engelm. ex C.L. Boynt. & Beadle R
Senecio quaylei T.M. Barkley TX (**Rev**) G1Q | known only from type
 localtion in Parker County; Turner et al. (2003) place under *S.*
ampullaceus Hook.
Shinnersoseris rostrata (A. Gray) Tomb s
Silphium albilorum A. Gray TX (**Rev**) R
Silphium integrifolium Michx. var. *laeve* Torr. & A. Gray
Solidago altiplanities C.E.S. Taylor & R. John Taylor OK, TX (**LEst**) R
Solidago capulinensis Cockerell & D.M. Andrews CO, NM (**Raton**) R
 G1 | Nesom & Lowrey 2011
Solidago gattingeri Chapm. ex A. Gray AR, MO, disj TN R G3?Q
Solidago mollis Bartl. var. *angustata* Shinners OK, TX (**Red**) R G5T3
Solidago speciosa Nutt. subsp. *pallida* Porter s
Symphotrichum eulae (Shinners) G.L. Nesom TX | known only from
 north-central Texas; recently collected in McLennan County
 (Hannick et al. 2013)
Symphotrichum fendleri (A. Gray) G.L. Nesom R
Symphotrichum praealtum (Poir.) G.L. Nesom var. *nebraskense*
 (Britton) G.L. Nesom IA, KS, NE
Symphotrichum sericeum (Vent.) G.L. Nesom
Townsendia texensis Larsen OK, TX (**LEst**) R
Verbesina lindheimeri B.L. Rob. & Greenm. TX (**Edw**) R
Vernonia guadalupensis A. Heller TX (**Edw**) RIP | possibly of hybrid
 origin involving *V. baldwinii* Torr. × *V. lindheimeri* Engelm. & A.
 Gray (= *V. × guadalupensis*)

Vernonia lindheimeri A. Gray & Engelm. TX, adj MX, disj AR R
Xanthisma spinulosum (Pursh) D.R. Morgan & R.L. Hartm. var. *glaberimum* (Rydb.) D.R. Morgan & R.L. Hartm.
Xanthisma texanum DC. var. *drummondii* (Torr. & A. Gray) A. Gray OK, TX

Berberidaceae

Berberis swaseyi Buckley TX (**Edw**) R G3

Boraginaceae

Lappula texana (Scheele) Britton var. *homosperma* A. Nelson & J.F. Macbr.

Lithospermum occidentale (Mack.) Weakly, Wistell & D. Estes
 [Onosmodium occidentale Mack.] R

Lithospermum subsetosum (Mack. & Bush) Weakly, Wistell & D. Estes
 [Onosmodium subsetosum Mack. & Bush] AR, MO, OK, disj TN R

Oreocarya cana A. Nelson [Cryptantha cana (A. Nelson) Payson] CO, MT, NE, SD, WY (**Niobr**) R

Brassicaceae

Cardamine carrii B.L. Turner TX (**Edw**) R | recently described (Turner 2012a) and probably of conservation concern

Dimorphocarpa candicans (Raf.) Rollins s

Erysimum asperum (Nutt.) DC.

Paysonia auriculata (Engelm. & A. Gray) O'Kane & Al-Shehbaz KS, OK, TX R

Physaria arenosa (Richardson) O'Kane & Al-Shehbaz subsp. *arenosa* R
Physaria arenosa (Richardson) O'Kane & Al-Shehbaz subsp. *argillosa* (Rollins & Shaw) O'Kane & Al-Shehbaz CO, NE, SD, WY R G5T3 | Beatty et al. 2003

Physaria bellii G.A. Mulligan CO R G2

Physaria brassicoides Rydb. R

Physaria calcicola (Rollins) O'Kane & Al-Shehbaz CO, NM (**Raton**) R G2

Physaria densiflora (A. Gray) O'Kane & Al-Shehbaz TX R

Physaria engelmannii (A. Gray) O'Kane & Al-Shehbaz OK, TX R

Physaria filiformis (Rollins) O'Kane & Al-Shehbaz AR, MO R G3/US-LT

Physaria gracilis (Hook.) O'Kane & Al-Shehbaz subsp. *nuttallii* (Torr. & A. Gray) O'Kane & Al-Shehbaz KS, OK, TX R

Physaria ovalifolia (Rydb.) O'Kane & Al-Shehbaz subsp. *alba* (Goodman) O'Kane & Al-Shehbaz KS, OK R G5T3?

Physaria ovalifolia (Rydb.) O'Kane & Al-Shehbaz subsp. *ovalifolia* R

Physaria recurvata (A. Gray) O'Kane & Al-Shehbaz TX R

Physaria sessilis (S. Watson) O'Kane & Al-Shehbaz TX (**Edw**) R

Selenia jonesii Cory TX (**Edw**) H G3

Streptanthus petiolaris A. Gray [Arabis petiolaris (A. Gray) A. Gray] TX (**Edw**) R

Thelypodium wrightii A. Gray subsp. *oklahomensis* Al-Shehbaz CO, OK (**Raton**) R

Cactaceae

Echinocereus milleri W. Blum, Kuenzler & Oldach TX (**Edw**) R G1

Echinocereus reichenbachii (Terscheck) Britton & Rose var. *baileyi* (Rose) N.P. Taylor OK, TX R G5T3

Echinocereus reichenbachii (Terscheck) Britton & Rose var. *perbellus* (Britton & Rose) L.D. Benson CO, NM, TX R

Echinocereus coccineus Engelm. subsp. *roemerii* (Muehlenp.) W. Blum, Mich. Lange & Rutow TX (**LUpl**) R

Opuntia mackensenii Rose var. *mackensenii* TX R | = *O. edwardsii* V.E. Grant & K.A. Grant (per Powell et al. 2008)

Sclerocactus brevihamatus (Engelm.) D.R. Hunt subsp. *tobuschii* (W.T. Marshall) N.P. Taylor TX (**Edw**) R G4T3/US-LE

Campanulaceae

Campanula reverchonii A. Gray TX (**LUpl**) R G2

Triodanis coloradoensis (Buckley) McVaugh TX (**Edw**), adj MX R

Triodanus leptocarpa (Nutt.) Nieuwl. s

Capparaceae

Cleomella angustifolia Torr s

Polanisia jamesii (Torr. & A. Gray) Iltis s

Caryophyllaceae

Minuartia michauxii (Fenzl) Farw. var. *texana* (B.L. Rob. ex Britton) Mattf. R

Paronychia depressa Nutt. ex Torr. & A. Gray CO, KS, NE, SD, WY (**Niobr**), disj NM R

Paronychia lindheimeri Engelm. & A. Gray TX (**Edw**), adj MX R

Paronychia virginica Spreng. var. *scoparia* (Small) Cory OK, TX, disj AR, MO R | Rohrer 1997

Convolvulaceae

Evolvulus arenarius R.T. Harms NM, TX s | recently described (Harms 2014) and possibly of conservation concern

Ipomoea costellata Torr. var. *edwardsiana* O'Kennon & G.L. Nesom TX (**Edw**) R G4T2

Ipomoea leptophylla Torr. s

Ipomoea shumardiana (Torr.) Shinners KS, OK, TX R G2

Crassulaceae

Sedum nuttallii Torr. & E. James ex Eaton [S. *nuttallianum* Raf.] R

Crossosomataceae

Forsellesia planiterium Ensign [Glossopetalon *planiterium* (Ensign) H. St. John] CO, NM, OK, TX (**LEst**) R

Forsellesia texensis Ensign [Glossopetalon *texense* (Ensign) H. St. John] TX (**Edw**) R G1

Cuscutaceae

Cuscuta plattensis A. Nelson NE, WY (**Niobr**), disj WA G1Q | Handley & Fertig 2002

Euphorbiaceae

Argythamnia aphoroides Müll. Arg. [Ditaxis *aphoroides* (Müll. Arg.) Pax] TX (**Edw**) R G2

Argythamnia humilis (Engelm. & A. Gray) Müll. Arg. var. *leiosperma* (Waterf.) OK, TX R

Argythamnia simulans J.W. Ingram [Ditaxis *simulans* (J.W. Ingram) Radcl.-Sm. & Govaerts] TX (**Edw**) R

Chamaesyce angusta (Engelm.) Small [Euphorbia *angusta* Engelm.] TX (**Edw**), adj MX R

Chamaesyce carunculata (Waterf.) Shinners KS, NM, OK, TX, adj MX s G3

Croton texensis Müll. Arg var. *texensis*

Euphorbia longicruris Scheele OK, TX, disj AR R

Euphorbia strictior Holz. NM, TX s G3

Stillingia texana I.M. Johnst. OK, TX, adj MX R

Tragia brevispica Engelm. & A. Gray TX R

Fabaceae

Amorpha canescens Pursh

Amorpha nana Nutt.

Amorpha roemeriana Scheele TX (**Edw**) R G3

Astragalus barrii Barneby MT, NE, SD, WY R G3 | Ladyman 2006a

Astragalus ceramicus Sheldon var. *filiformis* (A. Gray) F.J. Herm. s

Astragalus crassicaarpus Nutt. var. *berlanderi* Barneby TX (**Edw**) G5T3

Astragalus crassicaarpus Nutt. var. *trichocalyx* (Nutt.) Barneby R

Astragalus gracilis Nutt.

Astragalus hyalinus M.E. Jones CO, KS, MT, NE, WY (**Niobr**) R

Astragalus kentrophyta A. Gray var. *kentrophyta* R

Astragalus lindheimeri Engelm. ex A. Gray OK, TX R

Astragalus mollissimus Torr. var. *coryi* Tidest. TX (**Edw**) R G5T3

Astragalus mollissimus Torr. var. *mollissimus*

Astragalus plattensis Nutt. ex Torrey & A. Gray

Astragalus puniceus Osterh. var. *puniceus* CO, NM, OK (**Raton**), disj TX R G4T3?Q

Astragalus racemosus Pursh var. *racemosus*
Astragalus reflexus Torr. & A. Gray TX R G3
Astragalus sericoleucus A. Gray CO, KS, NE, WY (**Niobr**), disj NM R
Astragalus siliceus Barneby NM R G3
Astragalus wittmannii Barneby NM (**Raton**) R G3
Astragalus wrightii A. Gray TX R
Baptisia australis (L.) R. Br. var. *minor* (Lehm.) Fernald
Dalea arenicola (Wemple) B.L. Turner s
Dalea enneandra Nutt. R
Dalea hallii A. Gray TX (**Rev**) R G3
Dalea lanata Spreng. s | Turner (2006) extracted *D. austrotexana* B.L. Turner and *D. glaberrima* S. Watson from Barneby's (1977) broader circumscription of *D. lanata*
Dalea reverchonii (S. Watson) Shinners TX (**Rev**) R G2 | Taylor & O'Kennon 2013
Dalea sabinalis (S. Watson) Shinners TX (**Edw**) R GH | species has not been collected or reported since the 1950s and may be extinct (Poole et al. 2007)
Dalea tenuifolia (A. Gray) Shinners R
Dalea tenuis (J.M. Coult.) Shinners TX R
Dalea villosa (Nutt.) Spreng. s | *D. villosa* var. *grisea* (Torr. & A. Gray) of AR, LA, and east TX recognized by Turner et al. (2003) as *D. grisea* (Torr. & A. Gray) Shinners
Desmanthus leptolobus Torr. & A. Gray R
Desmodium illinoense A. Gray
Desmodium tweedyi Britton OK, TX R G3
Galactia texana (Scheele) A. Gray TX (**Edw**) R
Indigofera minuata Ortega var. *texana* (Buckley) B.L. Turner TX (**Lupl**) R
Lathyrus decaphyllus Pursh [*L. polymorphus* Nutt. (Broich 2009)] s [includes var. *incanus* (J.G. Sm. & Rydb.) Broich]
Lespedeza leptostachya Engelm. IA, IL, MN, WI G3/US-LT
Lespedeza texana Britton OK, TX, adj MX R
Lupinus plattensis S. Watson
Mimosa roemeriana Scheele OK, TX R
Mimosa rupertiana B.L. Turner s
Oxytropis lambertii Pursh var. *articulata* (Greene) Barneby R
Pediomelum argophyllum (Pursh) J.W. Grimes
Pediomelum cuspidatum (Pursh) Rydb. R
Pediomelum cyphocalyx (A. Gray) Rydb. TX R G3
Pediomelum digitatum (Nuttall ex Torrey & A. Gray) J.W. Grimes
Pediomelum esculentum (Pursh) Rydb.
Pediomelum hypogaeum (Nutt. ex Torr. & A. Gray) Rydb. var. *hypogaeum* R
Pediomelum hypogaeum (Nutt. ex Torr. & A. Gray) Rydb. var. *scapusum* (A. Gray) J.W. Grimes TX R
Pediomelum latestipulatum (Shinners) Mahler var. *appressum* (Ockendon) Gandhi & L.E. Br. TX (**Edw**) R
Pediomelum latestipulatum (Shinners) Mahler var. *latistipulatum* TX R
Pediomelum linearifolium (Torr. & A. Gray) J.W. Grimes R
Pediomelum reverchonii (S. Watson) Rydb. OK, TX (**Rev**) R G3
Pomaria brachycarpa (A. Gray) B.B. Simpson TX (**Edw**) R G2
Styphnolobium affine (Torr. & A. Gray) Walp. [*Sophora affinis* Torr. & A. Gray] OK, TX, disj AR, LA R

Fagaceae

Quercus havardii Rydb. NM, OK, TX s | AZ, NM, & UT populations formerly considered *Q. havardii* var. *tuckeri* S.L. Welsh now recognized as *Q. welschii* R.A. Denham

Fumariaceae

Corydalis curvisiliqua Engelm. subsp. *grandibracteata* (Fedde) G.B. Ownbey CO, KS, OK, TX

Gentianaceae

Centaurium beyrichii (Torr. & A. Gray) B.L. Rob. OK, TX R

Eustoma russellianum (Hook.) G. Don ex Sweet | Turner 2014
Frasera coloradensis (C.M. Rogers) D.M. Post CO (**Raton**) R G3 | Naumann 1991
Zeltnera texensis (Griseb.) G. Mans. ex J.S. Pringle [*Centaurium texense* (Griseb.) Fernald] AR, MO, OK, TX R

Lamiaceae

Brazoria enquistii M.W. Turner TX (**Lupl**) s G2
Hedeoma acinoides Scheele OK, TX R
Hedeoma reverchonii (A. Gray) A. Gray OK, TX R
Monarda clinopodioides A. Gray KS, OK, TX, disj LA s
Monarda punctata L. var. *intermedia* (E. L. McClint. & Epling) Waterf. TX s
Monarda stanfieldii Small TX (**Lupl**) s G3
Salvia engelmannii A. Gray TX R
Scutellaria bushii Britton AR, MO R G3
Scutellaria resinosa Torr. KS, OK, TX R
Warnockia scutellarioides (Engelm. & A. Gray) M.W. Turner OK, TX R

Linaceae

Linum berlandieri Hook.
Linum compactum A. Nelson
Linum rigidum Pursh

Loaceae

Eucnide bartonioides Zucc. var. *edwardsiana* B.L. Turner TX (**Edw**) R | recently described (Turner 2012b) and probably of conservation concern
Mentzelia chrysantha Engelm. ex K. Brandegee CO (**Ark**) R G2 | Anderson, D.G. 2006
Mentzelia nuda (Pursh) Torr. & A. Gray s
Mentzelia strictissima (Wootton & Standl.) J. Darl. NM, TX (**Mesc**) s

Lythraceae

Lythrum ovalifolium Koehn TX (**Edw**), adj MX RIP G3

Malvaceae

Callirhoe alcaeoides (Michx.) A. Gray
Callirhoe bushii Fernald AR, KS, MO, OK R G3
Callirhoe digitata Nutt. AR, MO, OK R
Callirhoe involucrata (Torr. & A. Gray) A. Gray var. *involucrata*
Callirhoe pedata (Nutt. ex Hook.) A. Gray AR, OK, TX R
Callirhoe scabruscula B.L. Rob. TX (**Red**) s G2/US-LE

Martyniaceae

Proboscidea sabulosa Correll NM, TX (**Mesc**), adj MX s G3

Nyctaginaceae

Mirabilis carletonii (Standl.) Standl. | recognized in some regional floras (GPFA 1986; Clark 1996; Kelso 2007) but placed in *M. glabra* (S. Watson) Standl. by Turner (1993) and in FNA (vol. 4, 2003)
Mirabilis exaltata (Standl.) Standl. KS, OK, TX G3 | recognized in some regional floras (GPFA 1986) but placed in *M. glabra* (S. Watson) Standl. by Turner (1993) and in FNA (vol. 4, 2003)
Mirabilis gigantea (Wootton & Standl.) Shinners TX (**Rev**) s
Mirabilis rotundifolia (Greene) Standl. CO (**Ark**) R G2

Onagraceae

Oenothera canescens Torr. & Frém. H
Oenothera capillifolia Scheele subsp. *capillifolia* [*Calylophus berlandieri* Spach subsp. *pinifolius* (Englemann) Towner] OK, TX, disj LA R
Oenothera cinerea (Wootton & Standl.) W. L. Wagner & Hoch subsp. *cinerea* [*Gaura villosa* Torr. subsp. *villosa*] s
Oenothera coloradensis (Rydb.) W.L. Wagner & Hoch [*Gaura neomexicana* Wootton subsp. *coloradensis* (Rydb.) P.H. Raven & D.P. Greg.] CO, NE, WY (**Niobr**) H G3T2/US-LT | Fertig 2000
Oenothera coryi W.L. Wagner TX (**Red**) R G3
Oenothera engelmannii (Small) Munz s

Oenothera glaucifolia W.L. Wagner & Hoch [*Stenosiphon linifolius* (Nutt. ex James) Heynh.] R
Oenothera harringtonii W.L. Wagner, Stockh. & W.M. Klein CO, NM (Raton) R G3 | Ladyman 2005
Oenothera macrocarpa Nutt. subsp. *fremontii* (S. Watson) W.L. Wagner KS, NE R GST3
Oenothera macrocarpa Nutt. subsp. *incana* (A. Gray) W.L. Wagner KS, NM, OK, TX (LEst) R GST3?
Oenothera macrocarpa Nutt. subsp. *macrocarpa* R
Oenothera macrocarpa Nutt. subsp. *oklahomensis* (Norton) W.L. Wagner KS, OK, TX (Red) R GST3?
Oenothera rhombipetala Nutt. ex Torr. & A. Gray s
Oenothera serrulata Nutt.
Oenothera suffulta (Engelm. & A. Gray) W.L. Wagner & Hoch subsp. *suffulta* [*Gaura suffulta* Engelm. & A. Gray] OK, TX
Oenothera triangulata (Buckley) W.L. Wagner & Hoch [*Gaura triangulata* Buckley] OK, TX (Red) R G3

Orobanchaceae

Agalinis densiflora (Benth.) S.F. Blake KS, OK, TX G3
Agalinis edwardsiana Pennell TX (Edw) R
Castilleja citrina Pennell KS, OK, TX R | Nesom & Egger 2014
Castilleja genevievana G.L. Nesom TX (Edw), adj MX R G3 | Nesom & Egger 2014
Castilleja purpurea (Nutt.) G. Don KS, OK, TX, disj MO R
Castilleja lindheimeri A. Gray [C. *purpurea* (Nutt.) G. Don var. *lindheimeri* (A. Gray) Shinners] TX (Edw) R GST3 | Nesom & Egger 2014

Papaveraceae

Argemone aurantiaca G.B. Ownbey TX (Edw) R
Argemone hispida A. Gray R
Argemone polyanthemus (Fedde) G.B. Ownbey s
Argemone squarrosa Greene subsp. *squarrosa* s

Plantaginaceae

Penstemon albidus Nutt.
Penstemon auriberbis Pennell CO, NM (Raton) R
Penstemon buckleyi Pennell s
Penstemon cobaea Nutt. R
Penstemon grandiflorus (Nutt.) s
Penstemon guadalupensis A. Heller TX (Edw) R G3
Penstemon helleri Small TX (Edw) R | sometimes placed in *P. triflorus* A. Heller (Diggs et al. 1999)
Penstemon haydenii S. Watson NE, disj WY s G1/US-LE | long thought to be endemic to the Nebraska Sandhills until outlying populations were discovered in central WY (Heidel 2012)
Penstemon oklahomensis Pennell OK, TX G3
Penstemon triflorus A. Heller TX (Edw) R G3
Penstemon versicolor Pennell CO (Ark) R G3?
Seymeria texana (A. Gray) Pennell TX (Edw) R G3

Polemoniaceae

Ipomopsis congesta (Hook.) V.E. Grant var. *pseudotypica* (Constance & Rollins) Dorn MT, SD, WY R GST3?
Phlox alyssifolia Greene R
Phlox andicola E.E. Nelson
Phlox drummondii Hook. subsp. *johnstonii* (Wherry) Wherry TX (Red) s GST3
Phlox drummondii Hook. subsp. *mcallesteri* (Whitehouse) Wherry TX s
Phlox oklahomensis Wherry KS, OK R G3 | Springer & Tyril 1989
Phlox pilosa L. subsp. *fulgida* Wherry
Phlox pilosa L. subsp. *longipilosa* (Waterf.) Locklear [*P. longipilosa* Waterf.] OK R G2Q
Phlox roemeriana Scheele TX R

Phlox villosissima (A. Gray) Small subsp. *latisejala* (Wherry) Locklear [*P. pilosa* L. subsp. *latisejala* Wherry]: TX (Rev), disj MX
Phlox villosissima (A. Gray) Small subsp. *villosissima* [*P. pilosa* L. subsp. *riparia* Wherry] TX (Edw) RIP | Locklear 2009, 2011

Polygonaceae

Eriogonum alatum Torr. var. *glabriusculum* Torr. NM, OK, TX (LEst) R
Eriogonum aliquantum Reveal NM (Raton) R G3
Eriogonum annuum Nutt. s
Eriogonum correllii Reveal OK, TX (LEst) R G2
Eriogonum helichrysoides (Gandoger) Prain in B.D. Jackson et al. [*E. effusum* Nutt. var. *rosmarinoides* Benth.] KS R G4G5T2?
Eriogonum jamesii Benth. var. *simplex* Gandoger KS R G5T2
Eriogonum nealleyi J.M. Coult. TX (Edw) R G2
Eriogonum nebraskense Rydb. CO, NE, WY (Niobr) R | possibly of hybrid origin involving *E. effusum* Nutt. × *E. pauciflorum* Pursh (= *E. x nebraskense*); also recognized as *E. pauciflorum* Pursh var. *nebraskense* (Rydb.) Reveal
Eriogonum pauciflorum Pursh var. *gnaphalodes* (Benth.) Reveal CO, NE, WY (Niobr) R
Eriogonum pauciflorum Pursh var. *pauciflorum* R
Eriogonum tenellum Torr. var. *ramosissimum* Benth. TX (LUpl) R GST3
Eriogonum visheri A. Nelson MT, ND, SD R G3 | Ladyman 2006c
Polygonum texense M.C. Johnst. TX H

Ranunculaceae

Delphinium carolinianum Walter var. *virescens* (Nutt.) R.E. Brooks
Delphinium treleasei Bush ex K.C. Davis AR, MO R G3

Rhamnaceae

Condalia hookeri M.C. Johnst. var. *edwardsiana* (Cory) M.C. Johnst. TX (Edw) R GST1Q

Rosaceae

Crataegus turnerorum Enquist TX (Edw) RIP G3Q
Prunus minutiflora Engelm. TX (Edw) R
Prunus pumila L. var. *besseyi* (L.H. Bailey) Waugh
Rosa foliolosa Nutt. ex Torr. & A. Gray

Rubiaceae

Galium texense A, Gray AR, OK, TX R
Hedyotis nigricans (Lam.) Fosberg var. *papillacea* B.L. Turner NM, TX (LEst) R GST3 | variety not recognized by Terrell (1986) who placed it in synonymy under *Stenaria nigricans* (Lam.) Terrell, but Turner considers it a distinct entity occurring primarily in the northern Texas panhandle (Turner 1995, 1997; Turner et al. 2003)

Scrophulariaceae

Besseya bullii (Eaton) Rydb. IA, IL, IN, MI, MN, OH, WI R G3

Solanaceae

Chamaesaracha darcy Averett OK, TX (Red) R
Chamaesaracha edwardsiana Averett TX (Edw) R

Valerianaceae

Valerianella amarella (Lindh. ex Engelm.) Krok R
Valerianella ozarkana Dyal AR, MO, OK R G3
Valerianella stenocarpa (Engelm.) Krok TX (Edw) RIP G3
Valerianella texana Dyal TX (LUpl) R G2

Violaceae

Viola pedatifida G. Don

Vitaceae

Parthenocissus heptaphylla (Buckley) Small TX (Edw)
Vitis acerifolia Raf.
Vitis cinerea (Engelm.) Millardet var. *helleri* (L.H. Bailey) M.O. Moore TX
Vitis monticola Buckley TX (Edw)

APPENDIX 2

CENTRAL GRASSLAND ENDEMIC PLANTS OF CONSERVATION CONCERN

The following taxa are plants of conservation concern based on NatureServe global conservation status rankings (G-rankings pertain to species, T-rankings to subspecific taxa). See Table 2 for more detailed descriptions of these rankings. Taxa also listed by the U. S. Fish and Wildlife Service as threatened or endangered are indicated by **LT** or **LE**. Taxa are arranged by center of endemism, if so associated, using the following abbreviations: Ark= Arkansas Valley Barrens; Edw = Edwards Plateau; LEst = Llano Estacado Escarpments; LUpl = Llano Uplift; Mesc = Mescalero-Monahans Dunes; Niobr = Niobrara-Platte Tablelands; Raton = Raton Tablelands; Red = Red Bed Plains; Rev = Reverchon Rocklands. State(s) and province(s) of occurrence are indicated.

Critically imperiled (G1/T1)

<i>Condalia hookeri</i>			
<i>var. edwardsiana</i>	Edw	TX	
<i>Echinocereus milleri</i>	Edw	TX	
<i>Elymus texensis</i>	Edw	TX	
<i>Forsellesia texensis</i>	Edw	TX	
<i>Perityle warnockii</i>	Edw	TX	
<i>Cuscuta plattensis</i>	Niobr	NE, WY	
<i>Solidago capulinensis</i>	Raton	CO, NM	
<i>Echinacea paradoxa</i>			
<i>var. neglecta</i>	Rev	OK	
<i>Senecio quaylei</i>	Rev	TX	
<i>Yucca necopina</i>	Rev	TX	
<i>Aschisma kansanum</i>		KS	
<i>Hymenoxys perpygaema</i>		TX	
<i>Molendoa ogalalensis</i>		KS, NE	
<i>Penstemon haydenii</i>		NE, WY	

Imperiled (G2/T2)

<i>Oonopsis puebloensis</i>	Ark	CO	
<i>Mentzelia chrysantha</i>	Ark	CO	
<i>Mirabilis rotundifolia</i>	Ark	CO	
<i>Argythamnia aphoroides</i>	Edw	TX	
<i>Eriogonum nealleyi</i>	Edw	TX	
<i>Ipomoea costellata</i>			
<i>var. edwardsiana</i>	Edw	TX	
<i>Pomaria brachycarpa</i>	Edw	TX	
<i>Eriogonum correllii</i>	LEst	OK, TX	
<i>Brazoria enquistii</i>	LUpl	TX	
<i>Campanula reverchonii</i>	LUpl	TX	
<i>Isoetes lithophila</i>	LUpl	TX	
<i>Packera texensis</i>	LUpl	TX	
<i>Tradescantia pedicellata</i>	LUpl	TX	
<i>Valerianella texana</i>	LUpl	TX	
<i>Cyperus onerosus</i>	Mesc	TX	
<i>Helianthus neglectus</i>	Mesc	NM, TX	
<i>Oenothera coloradensis</i>	Niobr	CO, NE, WY	
<i>Eurybia horrida</i>	Raton	CO, NM	
<i>Nolina greenei</i>	Raton	CO, NM, OK	
<i>Oonopsis monocephala</i>	Raton	CO	
<i>Packera spellenbergii</i>	Raton	NM	
<i>Physaria calcicola</i>	Raton	CO, NM	
<i>Callirhoe scabriuscula</i>	Red	TX	
<i>Dalea reverchonii</i>	Rev	TX	
<i>Asclepias meadii</i>	LT	IA, IL, IN, KS, MO, WI	
<i>Echinacea paradoxa</i>			
<i>var. paradoxa</i>		AR, MO	
<i>Eriogonum helichrysoides</i>		KS	
<i>Eriogonum jamesii</i>			
<i>var. simplex</i>		KS	

<i>Grindelia inornata</i>		CO, NM
<i>Ipomoea shumardiana</i>		KS, OK, TX
<i>Phlox pilosa</i>		
subsp. <i>longipilosa</i>		OK
<i>Physaria bellii</i>		CO

Vulnerable (G3/T3)

<i>Parthenium tetraeuris</i>	Ark	CO
<i>Penstemon versicolor</i>	Ark	CO
<i>Amorpha roemerana</i>	Edw	TX
<i>Astragalus crassicaarpus</i>		
<i>var. berlanderii</i>	Edw	TX
<i>Astragalus mollissimus</i>		
<i>var. coryi</i>	Edw	TX
<i>Berberis swaseyi</i>	Edw	TX
<i>Brickellia dentata</i>	Edw	TX
<i>Brickellia eupatorioides</i>		
<i>var. gracillima</i>	Edw	TX
<i>Castilleja genevievana</i>	Edw	TX
<i>Castilleja lindheimeri</i>	Edw	TX
<i>Crataegus turnerorum</i>	Edw	TX
<i>Hesperaloe parviflora</i>	Edw	TX
<i>Lythrum ovalifolium</i>	Edw	TX
<i>Penstemon guadalupensis</i>	Edw	TX
<i>Penstemon triflorus</i>	Edw	TX
<i>Perityle lindheimeri</i>		
<i>var. halmiifolia</i>	Edw	TX
<i>Perityle lindheimeri</i>		
<i>var. lindheimeri</i>	Edw	TX
<i>Sclerocactus breviphamatus</i>		
subsp. <i>tobuschii</i>	LEst	TX
<i>Selenia jonesii</i>	Edw	TX
<i>Seymeria texana</i>	Edw	TX
<i>Valerianella stenocarpa</i>	Edw	TX
<i>Yucca reverchonii</i>	Edw	TX
<i>Hedyotis nigricans</i>		
<i>var. papillacea</i>	LEst	NM, TX
<i>Oenothera macrocarpa</i>		
subsp. <i>incana</i>	LEst	KS, NM, OK, TX
<i>Eriogonum tenellum</i>		
<i>var. ramosissimum</i>	LUpl	TX
<i>Monarda stanfieldii</i>	LUpl	TX
<i>Ephedra coryi</i>	Mesc	NM, TX
<i>Eurytaenia hinckleyi</i>	Mesc	NM, TX
<i>Proboscidea sabulosa</i>	Mesc	NM, TX
<i>Parthenium alpinum</i>	Niobr	CO, NM, WY
<i>Astragalus puniceus</i>		
<i>var. puniceus</i>	Raton	CO, NM, OK, TX
<i>Astragalus wittmannii</i>	Raton	NM
<i>Eriogonum aliquantum</i>	Raton	NM

<i>Frasera coloradensis</i>	Raton	CO	<i>Eriogonum visheri</i>	MT, ND, SD
<i>Oenothera harringtonii</i>	Raton	CO, NM	<i>Euphorbia strictior</i>	NM, TX
<i>Oenothera coryi</i>	Red	TX	<i>Ipomopsis congesta</i>	
<i>Oenothera macrocarpa</i>			var. <i>pseudotypica</i>	MT, SD, WY
subsp. <i>oklahomensis</i>	Red	KS, OK, TX	<i>Lespedeza leptostachya</i> LT	IA, IL, MN, WI
<i>Oenothera triangulata</i>	Red	OK, TX	<i>Liatris glandulosa</i>	TX
<i>Phlox drummondii</i>			<i>Matelea edwardsensis</i>	TX
subsp. <i>johnstonii</i>	Red	TX	<i>Mirabilis exaltata</i>	KS, OK, TX
<i>Solidago mollis</i>			<i>Oenothera macrocarpa</i>	
var. <i>angustata</i>	Red	OK, TX	subsp. <i>fremontii</i>	KS, NE
<i>Dalea hallii</i>	Rev	TX	<i>Oonopsis engelmannii</i>	CO, KS
<i>Pediomelum reverchonii</i>	Rev	OK, TX	<i>Oonopsis foliosa</i>	CO
<i>Yucca pallida</i>	Rev	TX	<i>Pediomelum cyphocalyx</i>	TX
<i>Agalinis densiflora</i>		KS, OK, TX	<i>Penstemon oklahomensis</i>	OK, TX
<i>Ambrosia linearis</i>		CO	<i>Phlox oklahomensis</i>	KS, OK
<i>Astragalus barrii</i>		MT, NE, SD, WY	<i>Physaria arenosa</i>	
<i>Astragalus siliceous</i>		NM	subsp. <i>argillosa</i>	CO, NE, SD, WY
<i>Astragalus reflexus</i>		TX	<i>Physaria filiformis</i> LT	AR, MO
<i>Besseya bullii</i>		IA, IL, IN, MI, MN, OH, WI	<i>Physaria ovalifolia</i> subsp. <i>alba</i>	KS, OK
<i>Carex shinersii</i>		AR, KS, OK, TX	<i>Platanthera praeclara</i> LT	MB; IA, KS, MN, MO, ND,
<i>Chamaesyce carunculata</i>		KS, NM, OK, TX	NE, OK, SD	
<i>Chenopodium cycloides</i>		CO, KS, NE, NM, TX	<i>Scutellaria bushii</i>	AR, MO
<i>Delphinium treleasei</i>		AR, MO	<i>Solidago gatteringeri</i>	AR, MO
<i>Desmodium tweedyi</i>		OK, TX	<i>Valerianella ozarkana</i>	AR, MO, OK
<i>Dieteria canescens</i>			Possibly extinct (GH)	
var. <i>nebraskana</i>		NE, SD	<i>Dalea sabinalis</i>	Edw TX
<i>Echinacea atrorubens</i>		KS, OK, TX		
<i>Echinocereus reichenbachii</i>				
var. <i>baileyi</i>		OK, TX		

ACKNOWLEDGMENTS

This research would not have been possible without the body of floristic and ecological knowledge built up over the past half century by botanists working throughout the Central Grassland of North America. This paper is dedicated to the central figure of this scientific community, Ronald L. McGregor (1919–2012), who as director of the University of Kansas Herbarium (KANU; now the Ronald L. McGregor Herbarium) from 1954–1988 and catalyst and coordinator of the Great Plains Flora Association, helped bring into publication the *Atlas of the Flora of the Great Plains* (1977) and the *Flora of the Great Plains* (1986). Many thanks to Bruce W. Hoagland and Steven B. Rolfsmeier for manuscript review and suggested revisions.

REFERENCES

- ACKERFIELD, J. 2015. *Flora of Colorado*. Botanical Research Institute of Texas, Fort Worth, Texas, U.S.A.
- ALLISON, J.R. & T.E. STEVENS. 2001. Vascular flora of the Ketona Dolomite outcrops in Bibb County, Alabama. *Castanea* 66(1–2):154–205.
- AMOS, B.B. & C.M. ROWELL, JR. 1988. Floristic geography of woody and endemic plants. In: B.B. Amos & F.R. Gehlbach, eds. *Edwards Plateau vegetation: plant ecological studies in central Texas*. Baylor University Press, Waco, Texas, U.S.A. Pp. 25–42.
- ANDERSON, D.G. 2006. *Mentzelia chrysantha* Engelmann ex Brandegee (golden blazingstar): a technical conservation assessment. U.S.D.A. Forest Service, Rocky Mountain Region, Fort Collins, Colorado, U.S.A.
- ANDERSON, R.C. 2006. Evolution and origin of the Central Grassland of North America: Climate, fire, and mammalian grazers. *J. Torrey Bot. Soc.* 133(4):626–647.
- ANDERSON, R.C. & M.L. BOWLES. 1999. Deep-soil savannas and barrens in the Midwestern United States. In: R.C. Anderson, J.S. Fralish, & J.M. Baskin, eds. *Savannas, barrens, and rock outcrop communities of North America*, Cambridge University Press, New York, U.S.A. and Cambridge, U.K. Pp. 155–170.
- ARCHIBOLD, O.W. 1999. The aspen parkland of Canada. In: R.C. Anderson, J.S. Fralish, & J.M. Baskin, eds. *Savannas, barrens, and rock outcrop communities of North America*, Cambridge University Press, New York, U.S.A. and Cambridge, U.K. Pp. 406–420.
- ARMSTRONG, D.M. 1972. Distribution of mammals in Colorado. *Monogr. Mus. Nat. Hist., Univ. Kansas* 3:1–415.

- AXELROD, D.I. 1985. Rise of the grassland biome, central North America. *Bot. Rev.* 51(2):163–201.
- BARBER, S.C. 2008. A floristic study of the vascular plants of the Gypsum Hills and Redbed Plains area of southwestern Oklahoma. *Oklahoma Native Pl. Record* 8(1):4–36.
- BARE, J.E. & R.L. MCGREGOR. 1970. An introduction to the phytogeography of Kansas. *Univ. Kansas Sci. Bull.* 48(26):869–949.
- BARNEY, R.C. 1977. *Dalea* imagines. *Mem. New York Bot. Gard.* 27:1–892.
- BASKIN, J.M. & C.C. BASKIN. 1988. Endemism in rock outcrop plant communities of unglaciated eastern United States; an evaluation of the roles of edaphic, genetic and light factors. *J. Biogeogr.* 15:829–840.
- BASKIN, J.M. & C.C. BASKIN. 2000. Vegetation of limestone and dolomite glades in the Ozarks and Midwest regions of the United States. *Ann. Missouri Bot. Gard.* 87(2):286–294.
- BEATTY, B.L., W.F. JENNINGS, & R.C. RAWLINSON. 2003. *Lesquerella arenosa* (Richards.) Rydb. var. *argillosa* Rollins & Shaw (Great Plains bladderpod): A technical conservation assessment. U.S.D.A. Forest Service, Rocky Mountain Region, Fort Collins, Colorado, U.S.A.
- BIELMANN, A.P. & L.G. BRENNER. 1951. The recent intrusion of forests in the Ozarks. *Ann. Missouri Bot. Gard.* 38:261–282.
- BOLEN, E.G., L.M. SMITH, & H.L. SCHRAMM, JR. 1989. Playa lakes: Prairie wetlands of the southern High Plains. *BioScience* 39(9):615–623.
- BRENNAN, L.A. & W.P. KUVLESKY, JR. 2005. North American grassland birds: An unfolding conservation crisis? *J. Wildlife Managem.* 69(1):1–13.
- BRIDGES, E.L. & S.L. ORZELL. 1986. Distribution patterns of the non-endemic flora of Middle Tennessee limestone glades. *ASB Bull.* 33:155–166.
- BUCKALLEW, R.R. & G.M. CADDELL. Vascular flora of the University of Central Oklahoma Selman Living Laboratory, Woodward County, Oklahoma. *Proc. Oklahoma Acad. Sci.* 83:31–45.
- BUCHANAN, R., ed. 1984. *Kansas geology: An introduction to landscapes, rocks, minerals, and fossils.* University Press of Kansas, Lawrence, Kansas, U.S.A.
- BURGE, D.O. (& 13 others). 2016. Plant diversity and endemism in the California Floristic Province. *Madroño* 63(2):1–206.
- BUTHOD, A.K. & B.W. HOAGLAND. 2015. Contributions to the flora of Cimarron County and the Black Mesa area. *Oklahoma Native Pl. Record* 15:49–77.
- CADDELL, G.M. & K.D. RICE. 2012. Vascular flora of Alabaster Caverns State Park, Cimarron Gypsum Hills, Woodward County, Oklahoma. *Oklahoma Native Pl. Record* 12:43–62.
- CARIVEAU, A.B. & L. JOHNSON. 2007 Survey and assessment of playa wetlands in eastern Colorado; Neotropical migratory bird conservation act final report to the U.S. Fish and Wildlife Service, Rocky Mountain Bird Observatory, Brighton, Colorado, U.S.A.
- CARR, W.R. 2009. No place but Texas: An annotated list of plant taxa endemic to the Lone Star State (incomplete working draft). The Nature Conservancy of Texas, San Antonio, Texas, U.S.A.
- CARTWRIGHT, J.M. & W.J. WOLFE. 2016. Insular ecosystems of the southeastern United States: A regional synthesis to support biodiversity conservation in a changing climate. U.S. Geological Survey Professional Paper 1828.
- CHARBONEAU, J.L.M., B.E. NELSON, & R.L. HARTMAN. A floristic inventory of Phillips and Valley counties, Montana (U.S.A.). *J. Bot. Res. Inst. Texas* 7(2):847–878.
- CLARK, D.A. 1996. A floristic survey of the Mesa de Maya region, Las Animas, Colorado. *Natural History Inventory of Colorado No. 17.* University of Colorado Museum, Boulder, Colorado, U.S.A.
- CLEMENTS, F.E. 1920. *Plant indicators: The relation of plant communities to process and practice.* Carnegie Institution of Washington Publication 290. Washington, D.C., U.S.A.
- COCHRANE, T.S. & H.H. ILTIS. 2000. *Atlas of the Wisconsin prairie and savanna flora.* Department of Natural Resources Technical Bulletin No. 191, University of Wisconsin, Madison, Wisconsin, U.S.A.
- COLORADO NATURAL HERITAGE PROGRAM. 1997+. *Colorado Rare Plant Guide.* www.cnhp.colostate.edu. Latest update: June 30, 2014.
- COLORADO NATIVE PLANT SOCIETY. 1997. *Rare plants of Colorado*, 2nd ed. Falcon Press Publishing Company, Inc., Helena, Montana, U.S.A.
- COOPER, J.G. 1859. On the distribution of the forests and trees of North America, with notes on its physical geography. In: *Annual report to the Board of Regents of the Smithsonian Institution, showing the operations, expenditures, and conditions of the Institution for the year 1858.* Washington, D.C., U.S.A. Pp. 246–280.
- CORRELL, D.S. & M.C. JOHNSTON. 1970. *Manual of the vascular plants of Texas*, Texas Research Foundation, Renner, Texas, U.S.A.

- COSTEA, M., G.L. NESOM, & S. STEFANOVIC. 2006. Taxonomy of the *Cuscuta pentagona* complex (Convolvulaceae) in North America. *Sida* 22(1):151–175.
- CRIDLAND, A.A. 1959. The habitat of *Aschisma kansanum*. *Bryologist* 62:132–135.
- CURTIS, J.T. 1959. The vegetation of Wisconsin: An ordination of plant communities. University of Wisconsin Press, Madison, Wisconsin, U.S.A.
- CURTIS, N.M., Jr., W.E. HAM, & K.S. JOHNSON. 2008. Geomorphic provinces of Oklahoma. In: K.S. Johnson & K.V. Luza, eds, Earth sciences and mineral resources of Oklahoma. Oklahoma Geological Survey Educational Publication 9, University of Oklahoma, Norman, Oklahoma, U.S.A. Pp. 8.
- DECKER, K. 2006. *Asclepias uncialis* Greene (wheel milkweed): A technical conservation assessment. U.S.D.A. Forest Service, Rocky Mountain Region, Fort Collins, Colorado, U.S.A.
- DECKER, K. 2007. The potential distribution and landscape integrity of *Frankenia jamesii* (James' seaheath) in southeastern Colorado. Colorado Natural Heritage Program, Colorado State University, Fort Collins, Colorado, U.S.A.
- DIAMOND, D.D. & F.E. SMEINS. 1985. Composition, classification and species response patterns of remnant tallgrass prairies in Texas. *Amer. Midl. Naturalist* 113(2):294–308.
- DIAMOND, D.D. & F.E. SMEINS. 1988. Gradient analysis of remnant True and Upper Coastal Prairie grasslands of North America. *Canad. J. Bot.* 66:2152–2161.
- DIAMOND, D.D. & F.E. SMEINS. 1993. The native plant communities of the Blackland Prairie. In: M.R. Sharpless & J.C. Yelderman, Jr., eds. The Texas Blackland Prairie, land, history, and culture. Baylor University Program for Regional Studies, Waco, Texas, U.S.A. Pp. 66–81.
- DICK-PEDDIE, W.A. 1993. New Mexico vegetation: Past, present, and future. University of New Mexico Press, Albuquerque, New Mexico, U.S.A.
- DIFFENDAL, R.F., Jr. 1991. Plate tectonics, space, geologic time, and the Great Plains: A primer for non-geologists. *Great Plains Quart.* 11(2):83–102.
- DIFFENDAL, R.F., Jr. 2017. Great Plains geology. University of Nebraska Press, Lincoln, Nebraska, U.S.A.
- DIGGS, G.M. & B.L. LIPSCOMB. 2014. The ferns and lycophytes of Texas. Botanical Research Institute of Texas, Fort Worth, Texas, U.S.A.
- DIGGS, G.M., Jr., B.L. LIPSCOMB, & R.J. O'KENNON. 1999. Shinners & Mahler's illustrated flora of North Central Texas. *Sida, Bot. Misc.* 16.
- DIGGS, G.M., Jr., B.L. LIPSCOMB, M.D. REED, & R.J. O'KENNON. 2006. Illustrated flora of East Texas, vol. 1. *Sida, Bot. Misc.* 26.
- DHILLION, S.S. & M.H. MILLS. 1999. The sand shinnery oak (*Quercus havardii*) communities of the Llano Estacado: History, structure, ecology, and restoration. In: R.C. Anderson, J.S. Fralish, & J.M. Baskin, eds. Savannas, barrens, and rock outcrop communities of North America, Cambridge University Press, New York, U.S.A. and Cambridge, U.K. Pp. 262–274.
- DORN, R.D. 2001. Vascular plants of Wyoming, 3rd ed. Mountain West Publishing, Cheyenne, Wyoming, U.S.A.
- DORR, L.J. 1990. Revision of the North American genus *Callirhoe* (Malvaceae). *Mem. New York Bot. Gard.* 56:1–76.
- DYKSTERHUIS, E.J. 1946. The vegetation of the Fort Worth Prairie. *Ecol. Monogr.* 16:1–29.
- ESCUADERO, A., S. PALACIO, F.T. MAESTRE, & A.L. LUZURIAGA. 2014. Plant life on gypsum: A review of its multiple facets. *Biol. Rev.* 90:1–18.
- ESTILL, J.C. & M.B. CRUZAN. 2001. Phytogeography of rare plant species endemic to the southeastern United States. *Castanea* 66(1–2):3–23.
- EVANS, C.S. 2010. Playas in Kansas and the High Plains. Kansas Geological Survey Public Information Circular 30. Kansas Geological Survey, University of Kansas, Lawrence, Kansas, U.S.A.
- FENNEMAN, N.M. 1931. Physiography of the western United States. McGraw-Hill, New York, New York, U.S.A.
- FERTIG, W. 2000. Status review of the Colorado butterfly plant (*Gaura neomexicana* ssp. *coloradensis*). Unpublished report prepared for the Wyoming Cooperative Fish and wildlife Research Unit, U.S. Fish and Wildlife Service, and Wyoming Game and Fish Department by the Wyoming Natural Diversity Database, Laramie, Wyoming, U.S.A.
- FLORA OF NORTH AMERICA EDITORIAL COMMITTEE, EDs. [FNAEC] 1993–. Flora of North America north of Mexico. 18+ vols. Oxford University Press, New York, U.S.A. and Oxford, U.K.
- FLORES, D. 2010. Caprock Canyonlands: Journeys into the heart of the southern plains, 20th anniversary ed. Texas A. & M. University Press, College Station, Texas, U.S.A.
- FOTI, T.L. & G.A. BUKENHOFER. 1998. A description of the sections and subsections of the Interior Highlands of Arkansas and Oklahoma. *J. Arkansas Acad. Sci.* 52:53–62.
- FOTI, T.L., S. SIMON, D. ZOLLNER, & M. HATTENBACH. 2003. Blackland prairie landscapes of southwestern Arkansas: Historical

- perspective, present status, and restoration potential. In: E. Peacock & T. Schauwecker, eds. Blackland prairies of the Gulf Coastal Plain. University of Alabama Press, Tuscaloosa, Alabama, U.S.A. Pp. 94–109.
- FOTI, T. & C.T. WITSELL. 2013. Effects of physical factors on the distribution of native flora and vegetation in the Natural Divisions of Arkansas. In: J.L. Gentry, G.P. Johnson, B.T. Baker, C.T. Witsell, & J.D. Ogle, eds. Atlas of the vascular plants of Arkansas, Arkansas Vascular Flora Committee, Fayetteville, Arkansas, U.S.A. Pp. 17–40.
- FOWLER, N.L. & D.W. DUNLAP. 1986. Grassland vegetation of the eastern Edwards Plateau. *Amer. Midl. Nat.* 115(1):146–155.
- GAGE, A.M., S.K. OLIMB, & J. NELSON. 2016. Plowprint: Tracking cumulative cropland expansion to target grassland conservation. *Great Plains Res.* 26(2):107–116.
- GENTRY, J.L., G.P. JOHNSON, B.T. BAKER, C.T. WITSELL, & J.D. OGLE, eds. 2013. Atlas of the vascular plants of Arkansas, Arkansas Vascular Flora Committee, Fayetteville, Arkansas, U.S.A.
- GRAVES, G.R. 2016. First record of *Clematis fremontii* S. Watson from Arkansas. *Castanea* 81(4):333–335.
- GREAT PLAINS FLORA ASSOCIATION [GPFA]. 1977. Atlas of the flora of the Great Plains. Iowa State University Press, Ames, Iowa, U.S.A.
- GREAT PLAINS FLORA ASSOCIATION [GPFA]. 1986. Flora of the Great Plains. University Press of Kansas, Lawrence, Kansas, U.S.A.
- GRIFFITH, G.E., S.A. BRYCE, J.M. OMERNIK, J.A. COMSTOCK, A.C. ROGERS, B. HARRISON, S.L. HATCH, & D. BEZANSON. 2004. Ecoregions of Texas, U.S. Environmental Protection Agency, Corvallis, Oregon, U.S.A., 1:2,500,000.
- HANBERRY, B.B., D.T. JONES-FARRAAND, & J.M. KABRICK. 2014. Historical open forest ecosystems in the Missouri Ozarks: Reconstruction and restoration targets. *Ecol. Res.* 32(4):407–416.
- HANDLEY, J & W. FERTIG. 2002. *Cuscuta plattensis* Wyoming dodder. State species abstract, Wyoming Natural Diversity Database, University of Wyoming, Laramie, Wyoming, U.S.A.
- HARDY, J.P. 1991. The vascular flora of Banner County, Nebraska. *Trans. Nebraska Acad. Sci.* 18:109–126.
- HARMS, R. T. 2014. A new species of *Evolvulus* (Convolvulaceae) from the High Plains of the Texas/New Mexico border. *Phytoneuron* 2014-20:1–20.
- HATTIN, D.E. 1975. Stratigraphy and depositional environments of Greenhorn Limestone (Upper Cretaceous) of Kansas. *Kansas Geol. Survey Bull.* 209.
- HATTIN, D.E. 1982. Stratigraphy and depositional environment of Smoky Hill Chalk Member, Niobrara Chalk (Upper Cretaceous) of the type area, western Kansas. *Kansas Geol. Survey Bull.* 225.
- HAUKOS, D.A. & L.M. SMITH. 1997. Common flora of the playa lakes. Texas Tech University Press, Lubbock, Texas, U.S.A.
- HAZLETT, D.L. 1998. Vascular plant species of the Pawnee National Grassland. Gen. Tech. Rep. RMRS-GTR-17, U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station, Fort Collins, Colorado, U.S.A.
- HAZLETT, D.L. 2004. Vascular plant species of the Comanche National Grasslands in southeastern Colorado. Gen. Tech. Rep. RMRS-GTR-130. U.S. Department of Agriculture Forest Service, Rocky Mountain Research Station, Fort Collins, Colorado, U.S.A.
- HAZLETT, D.L., M.H. SCHIEBOUT, & P.L. FORD. 2009. Vascular plants and a brief history of the Kiowa and Rita Blanca National Grasslands. Gen. Tech. Rep. RMRSGTR-233. U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station, Fort Collins, Colorado, U.S.A.
- HEIDEL, B. 2012. Status of *Penstemon haydenii* (Blowout Penstemon) in Wyoming. 2012. Prepared for Bureau of Land Management—Rawlins and Rock Springs Field Offices and Wyoming State Office. Wyoming Natural Diversity Database, University of Wyoming, Laramie, Wyoming, U.S.A.
- HEIDEL, B. & J. HANDLEY. 2004. *Parthenium alpinum* (Nutt.) Torr. & Gray (alpine feverfew): A technical conservation assessment. U.S.D.A. Forest Service, Rocky Mountain Region, Fort Collins, Colorado, U.S.A.
- HEIL, K.D., S.L. O'KANE, JR., L.M. REEVES, & A. CLIFFORD. 2013. Flora of the Four Corners region: Vascular plants of the San Juan River drainage, Arizona, Colorado, New Mexico, and Utah. Missouri Botanical Garden Press, St. Louis, Missouri, U.S.A.
- HENWOOD, W.D. 2010. Toward a strategy for the conservation and protection of the world's temperate grasslands. *Great Plains Res.* 20(1):121–134.
- HILL, R.T. 1901. Geography and geology of the Black and Grand Prairies, Texas. U.S. Geological Survey. Annual Report 21. United States Geological Survey, Washington, D.C., U.S.A.
- HOAGLAND, B.W. 2000. The vegetation of Oklahoma: A classification of landscape mapping and conservation planning. *Southw. Nat.* 45:385–420.
- HOAGLAND, B.W. & A.K. BUTHOD. 2005. Vascular flora of a gypsum dominated site in Major County, Oklahoma. *Proc. Okla. Acad. Sci.* 85:1–8.
- HOAGLAND, B.W., I.H. BUTLER, F.L. JOHNSON, & S. Glenn. 1999. The Cross Timbers. In: R.C. Anderson, J.S. Fralish, & J.M. Baskin,

- eds. Savannas, barrens, and rock outcrop communities of North America, Cambridge University Press, New York, U.S.A. and Cambridge, U.K. Pp. 231–245.
- HOAGLAND, B.W. & S.L. COLLINS. 1997. Heterogeneity in shortgrass prairie vegetation; the role of playa lakes. *J. Veg. Sci.* 8:277–286.
- HOAGLAND, B.W. & F.L. JOHNSON. 2001. Vascular flora of the Chickasaw National Recreation Area, Murray County, Oklahoma. *Castanea* 66(4):383–400.
- HOEKSTRA, J.M., T.M. BOUCHER, T.H. RICKETTS, & C. ROBERTS. 2005. Confronting a biome crisis: Global disparities of habitat loss and protection. *Ecol. Lett.* 8:23–29.
- HOOD, H.C. & J.R. UNDERWOOD, JR. 2001. Geology of Palo Duro Canyon. In: D.F. Guy, ed. *The story of Palo Duro Canyon*. Texas Tech University Press, Lubbock, Texas, U.S.A. Pp. 3–34.
- HOLLIDAY, V.T. 2001. Stratigraphy and geochronology of upper Quaternary eolian sand on the Southern High Plains of Texas and New Mexico, United States. *GSA Bull.* 113(1):88–108.
- HORN, D.D. & J. Shaw. 2007. Noteworthy collections: Tennessee. *Castanea* 72(1):48–49.
- HUNT, C.B. 1974. *Natural regions of the United States and Canada*. W.H. Freeman, San Francisco, California, U.S.A.
- JERCINOVIC, E. 2007. The status of genus *Chamaesyce* in New Mexico. *New Mexico Botanist* 40:1–14.
- JOHNSON, A.F., W.W. BAKER, L.C. ANDERSON, & A.K. GHOLSON, JR. 2013. Flora of calcareous upland glades in Gadsden and Jackson counties, Florida. *J. Bot. Res. Inst. Texas* 7(1):475–494.
- JOHNSTON, M.C. 1969. *Polygonum texense* M.C. Johnston (Polygonaceae), new species from South Plains and Edwards Plateau. *Southw. Nat.* 14(2):257–258.
- KARTESZ, J.T. 2015. North American Plant Atlas. (<http://bonap.net/napa>). Chapel Hill, N.C. [maps generated from Kartesz, J.T. 2015. Floristic Synthesis of North America, Version 1.0. Biota of North America Program (BONAP). (in press)]
- KAUL, R.B., D. SUTHERLAND, & S. ROLFSMEIER. 2011. *The flora of Nebraska*, 2nd ed. School of Natural Resources, University of Nebraska-Lincoln, Lincoln, Nebraska, U.S.A.
- KELSO, S., C. HALL, & G. MAENTZ. 2001. The role of landscape anomalies in regional plant conservation. In: J. Maschinski & L. Holter, eds. *Southwestern rare and endangered plants: Proceedings of the third conference; 2000 September 25–28; Flagstaff, Arizona*. RMRS-P-23. U.S. Department of Agriculture, Forest Service, Fort Collins, Colorado, U.S.A. Pp. 13–19.
- KELSO, S., K. HECKMAN, J. LAWTON, & G. MAENTZ. 1996. Endemic calciphiles of the Middle Arkansas Valley, Colorado. In: J. Maschinski, H.D. Hammond, & L. Holter, eds. *Southwestern rare and endangered plants: Proceedings of the second conference; 1995 September 11–14, Flagstaff, Arizona*. Gen. Tech. Rep. RM-GTR-283, U.S. Department of Agriculture, Forest Service, Fort Collins, Colorado, U.S.A. Pp. 270–280.
- KELSO, S., N. BOWER, K.E. HECKMANN, P.M. BEARDSLEY, & D.G. GREVE. 2003. Geobotany of the Niobrara Chalk Barrens in Colorado: A study of edaphic endemism. *W. N. Amer. Naturalist* 63(3):299–313.
- KELSO, T., N. BOWER, P. HALTEMAN, K. TENNEY, & S. WEAVER. 2007. Dune communities of SE Colorado: Patterns of rarity, disjunction and succession. In: P. Barlow-Irlick, J. Anderson, C. McDonald, eds. *Southwestern rare and endangered plants: Proceedings of the fourth conference; 2004 March 22–26, Las Cruces, New Mexico*. RMRS-P-48CD, U.S. Department of Agriculture, Forest Service, Fort Collins, Colorado, U.S.A. Pp. 39–48.
- KNIGHT, D.H. 1994. *Mountains and plains: The ecology of Wyoming landscapes*. Yale University Press, New Haven, Connecticut, U.S.A.
- KNIGHT, D.H. 1999. Ponderosa and limber pine woodlands. In: R.C. Anderson, J.S. Fralish, & J.M. Baskin, eds. *Savannas, barrens, and rock outcrop communities of North America*, Cambridge University Press, New York, U.S.A. and Cambridge, U.K. Pp. 249–261.
- KRAFT, N.J.B., B.G. BALDWIN, & D.D. ACKERLY. 2010. Range size, taxon age and hotspots of neoendemism in the California flora. *Diversity & Distrib.* 16:403–413.
- KRUCKEBERG, A.R. 1986. An essay: The stimulus of unusual geologies for plant speciation. *Syst. Bot.* 11:455–463.
- KRUCKEBERG, A.R. 2002. *Geology and plant life: The effects of landforms and rock types on plants*. University of Washington Press, Seattle, Washington, U.S.A.
- KRUCKEBERG, A.R. & D. RABINOWITZ. 1985. Biological aspects of endemism in higher plants. *Ann. Rev. Ecol. Syst.* 16:447–479.
- KUCHLER, A.W. 1985. *Potential natural vegetation of the conterminous United States* [map], revised. American Geographical Society, New York, New York, U.S.A.
- KUHN, B., B.E. NELSON, & R.L. HARTMAN. 2011. A floristic inventory of the Cimarron National Grassland (Kansas) and the Comanche National Grassland (Colorado). *J. Bot. Res. Inst. Texas* 5(2):753–772.
- LADYMAN, J.A.R. 2005. *Oenothera harringtonii* Wagner, Stockhouse & Klein (Colorado Springs evening primrose): A technical conservation assessment. U.S.D.A. Forest Service, Rocky Mountain Region, Fort Collins, Colorado, U.S.A.

- LADYMAN, J.A.R. 2006a. *Astragalus barrii* Barneby (Barr's milkvetch): A technical conservation assessment. U.S.D.A. Forest Service, Rocky Mountain Region, Fort Collins, Colorado, U.S.A.
- LADYMAN, J.A.R. 2006b. *Chenopodium cycloides* A. Nelson (sandhill goosefoot): A technical conservation assessment. U.S.D.A. Forest Service, Rocky Mountain Region, Fort Collins, Colorado, U.S.A.
- LADYMAN, J.A.R. 2006c. *Eriogonum visleri* A. Nelson (Visher's buckwheat): A technical conservation assessment. U.S.D.A. Forest Service, Rocky Mountain Region, Fort Collins, Colorado, U.S.A.
- LAUENROTH, W.K., I.C. BURKE, & M.P. GUTMANN. 1999. The structure and function of ecosystems in the central North American grassland region. *Great Plains Res.* 9(2):223–259.
- LAVIN, M. & C. SEIBERT. 2011. Great Plains flora? Plant geography of eastern Montana's lower elevation shrub-grass dominated vegetation. In: C.L. Wambolt et al., comps. 15th wildland shrub symposium – Shrublands: Wildlands and wildlife habitats, 2008 June 17–19, Bozeman, Montana. *Natural Resources and Environmental Issues* vol. 16, article 1, Utah State University, Logan, Utah, U.S.A.
- LAWLESS, P.J., J.M. BASKIN, & C.C. BASKIN. 2006. Xeric limestone prairies of Eastern United States: Review and synthesis. *Bot. Rev.* 72(3):235–272.
- LEE, W.T. 1902. The canyons of southeastern Colorado. *J. Geogr.* 1(8):357–370.
- LEE, W.T. 1903. The canyons of northeastern New Mexico. *J. Geogr.* 2(2):63–82.
- LEE, W.T. 1912. Extinct volcanoes of northeast New Mexico. *Amer. Forest.* 18(6):357–365.
- LESICA, P. 2012. Manual of Montana vascular plants. Botanical Research Institute of Texas, Fort Worth, Texas, U.S.A.
- LLADO, L. & M.C. SLATTERLY. 2015. Fine scale temporal variations of surface moisture in topographically controlled Muhly grass seeps. *Ecology* 96(3):371–381.
- LOCKLEAR, J.H. 1989. The status of *Ambrosia linearis* (Rydb.) Payne in Colorado. Report to the U.S. Fish and Wildlife Service Endangered Species Office, Denver, Colorado, U.S.A.
- LOCKLEAR, J.H. 1990. A Colorado specialty: *Ambrosia linearis*. *Aquilegia* [Colorado Native Pl. Soc. News.] 14(5):10–11.
- LOCKLEAR, J.H. 2011. *Phlox*: A natural history and gardener's guide. Timber Press, Portland, Oregon, U.S.A.
- LOCKLEAR, J.H. 2014. Reconnaissance survey of rock outcrop communities in the Kimball Grasslands of Nebraska. Prepared for the Nebraska Natural Heritage Program, Nebraska Game and Parks Commission, Lincoln, Nebraska, U.S.A.
- LONGING, S., S. DISCUA, & J. COKENDOLPHER. 2014. Surveys and habitat assessment of endemic insects at the Monahans dune system. Final report prepared for the Texas Parks and Wildlife Department, Texas Tech University, Lubbock, Texas, U.S.A.
- LOOMAN, J. & K.F. BEST. 1987. Budd's flora of the Canadian Prairie Provinces. Research Branch Agriculture Canada Publication 1662.
- MACROBERTS, B.R., M.H. MACROBERTS, C.S. REID, & P.L. FAULKER, 2009. Vascular flora of Morse Clay prairies in northwestern Louisiana. *J. Bot. Res. Inst. Texas* 3(1):355–366.
- MACROBERTS, B.R., M.H. MACROBERTS, & C.T. WITSELL. 2011. Small-scale vascular plant species richness in southwestern Arkansas blackland prairies. *J. Bot. Res. Inst. Texas* 5(2):743–751.
- MACROBERTS, M.H., B.R. MACROBERTS, & L.S. JACKSON. 2003. Louisiana prairies. In: E. Peacock & T. Schauwecker, eds. *Blackland prairies of the Gulf Coastal Plain*. University of Alabama Press, Tuscaloosa, Alabama, U.S.A. Pp. 80–93.
- MACROBERTS, M.H. B.R. MACROBERTS, B.A. SORRIE, & R.E. EVANS. 2002. Endemism in the West Gulf Coastal Plain; importance of xeric habitats. *Sida* 20(2):767–780.
- MAHLER, H.D., Jr., G.F. ENGELMANN, & R.D. SHUSTER. 2003. *Roadside geology of Nebraska*. Mountain Press Publishing Company, Missoula, Montana, U.S.A.
- McMILLAN, M.E., C.L. ANGEVINE, & P.L. HELLER. 2002. Postdepositional tilt of the Miocene-Pliocene Ogallala group on the western Great Plains: Evidence of late Cenozoic uplift of the Rocky Mountains. *Geology* 30(1):63–66.
- MENGES, E.S. 1999. Ecology and conservation of Florida Scrub. In: R.C. Anderson, J.S. Fralish, & J.M. Baskin, eds. *Savannas, barrens, and rock outcrop communities of North America*, Cambridge University Press, New York, U.S.A. and Cambridge, U.K. Pp. 7–22.
- MINK, J.N., J.R. SINGHURST, & W.C. HOLMES. 2012. A new species of *Hymenoxys* (Asteraceae, Helenieae, Tetraneuridinae) from Texas. *Novon* 22:56–59.
- MOHR, C. 1901. Plant life of Alabama. *Contr. U.S. Natl. Herb.* 6:1–921.
- MOIR, W.H. 1969. Steppe communities in the foothills of the Colorado Front Range and their relative productivities. *Amer. Midl. Naturalist* 81:331–340.

- MUEGLER, W.F. & W.L. STEWART. 1980. Grassland and shrubland habitat types of western Montana. Gen. Tech. Rep Int-66, U.S. Department of Agriculture, Forest Service, Intermountain Forest and Range Experimental Station, Ogden, Utah, U.S.A.
- MUHS, D.R. & V.T. HOLLIDAY. 1995. Evidence of active dune sand on the Great Plains in the 19th century from accounts of early explorers. *Quaternary Res.* 43:198–208.
- MUHS, D.R. & V.T. HOLLIDAY. 2001. Origin of late Quaternary dune fields on the Southern High Plains of Texas and New Mexico. *Bull. Geol. Soc. Amer.* 113(1):75–87.
- NAUMANN, T.S. 1991. Status report for *Frasera coloradensis*. Unpublished report prepared for the Colorado Natural Areas Program, Denver, Colorado, U.S.A.
- NEELY, B., S. PANJABI, & J. HANDWERK. 2011. Arkansas Valley Barrens: Conservation action plan 2011 update. The Nature Conservancy and the Colorado Natural Heritage Program, Colorado State University, Fort Collins, Colorado, U.S.A.
- NEID, S., K. DECKER, J. HANDWERK, & S.S. PANJABI. 2007. Rare plant surveys on the Piñon Canyon Maneuver Site 2006–2007. Colorado Natural Heritage Program, Colorado State University, Fort Collins, Colorado, U.S.A.
- NEID, S. & J. HANDWERK. 2007. Rare plant surveys on Fort Carson 2006–2007. Colorado Natural Heritage Program, Colorado State University, Fort Collins, Colorado, U.S.A.
- NELSON, A. 1899. New plants from Wyoming VI. *Bull. Torrey Bot. Club* 26:122–134.
- NELSON, P.W. 2005. The terrestrial natural communities of Missouri, revised ed. The Missouri Natural Areas Committee, Jefferson City, Missouri, U.S.A.
- NELSON, P. & D. LADD. 1980. Preliminary report on the identification, distribution, and classification of Missouri glades. In: C. Kucera, ed. Proceedings of the seventh North American Prairie conference, Southwest Missouri State University, Springfield, Missouri, U.S.A. Pp. 59–76.
- NESOM, G.L. & T.K. LOWREY. 2011. *Solidago capulinensis* (Asteraceae: Astereae) redividus. *Phytoneuron* 2011-24:1–22.
- NESOM, G.L. & R.J. O'KENNON. 2001. Two new species of *Liatris* series *Punctatae* (Asteraceae: Eupatorieae) centered in North Central Texas. *Sida* 19:767–787.
- NESOM, G.L. & R. J. O'KENNON. 2008. Major plant communities of Lake Meredith National Recreational Area and Alibates Flint Quarries National Monument. *Phytologia* 90(3):391–401.
- NEW MEXICO RARE PLANT TECHNICAL COUNCIL. 1999. New Mexico Rare Plants. Albuquerque, NM: New Mexico Rare Plants Home Page. <http://nmrareplants.unm.edu> (Latest update: 21 July 2016).
- NOSS, R.F. 2013. *Forgotten grasslands of the South: Natural history and conservation*. Island Press, Washington, D.C., U.S.A.
- NOSS, R.F., W.J. PLATT, B.A. SORRIE, A.S. WEAKLEY, D.B. MEANS, J. COSTANZA, & R.K. PEET. 2014. How global biodiversity hotspots may go unrecognized; lessons from the North American Coastal Plain. *Diversity Distrib.* 21:236–244.
- NUZZO, V. 1986. Extent and status of Midwest oak savanna: Presettlement and 1985. *Natural Areas J.* 6:6–36.
- O'KENNON, R.J. & K.N. TAYLOR. 2015. *Cyperus granitophilus* (Cyperaceae), a granite outcrop endemic, new for Texas and Oklahoma (U.S.A.). *J. Bot. Res. Inst. Texas* 9(1):251–257.
- PETERSON, R.S. & C.S. BOYD. 1998. Ecology and management of sand shinnery communities: A literature review. Gen. Tech. Rep RMRS-GTR-16, U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station, Fort Collins, Colorado, U.S.A.
- POOLE, J.M. W.R. CARR, D.M. PRICE, & J.R. SINGHURST. 2007. *Rare plants of Texas*. Texas A & M University Press, College Station, Texas, U.S.A.
- PRINGLE, J.S. & T. WITSELL. 2005. A new species of *Sabatia* (Gentianaceae) from Saline County, Arkansas. *Sida* 21(3):1249–1262.
- RADUSKI, A.R., L.H. RIESEBERG, & J.L. STRASBURG. 2010. Effective population size, gene flow, and species status in a narrow endemic sunflower, *Helianthus neglectus*, compared to its widespread sister species, *H. petiolaris*. *Int. J. Mol. Sci.* 11:492–506.
- RAMALEY, F. 1939. Sand-hill vegetation of northeastern Colorado. *Ecol. Monogr.* 9(1):1–51.
- RAVIN, P.H., & D.I. AXELROD. 1978 (reprinted 1995). *Origins and relationships of the California flora*. California Native Plant Society, Sacramento, California, U.S.A.
- REEVES, C.C., JR. 1972. Tertiary-Quaternary stratigraphy and geomorphology of West Texas and southeastern New Mexico. In: V.C. Kelly & F.D. Trauger, eds. *East-central New Mexico*, New Mexico Geological Survey 23rd annual fall field conference guidebook. New Mexico Geological Society, Socorro, New Mexico, U.S.A. Pp. 108–117.
- REID, C.S., P.L. FAULKNER, M.H. MACROBERTS, B.R. MACROBERTS, & M. BORDELON. 2010. Vascular flora and edaphic characteristics of saline prairies in Louisiana. *J. Bot. Res. Inst. Texas* 4(1):357–379.

- RISKIND, D.H. & D.D. DIAMOND. 1988. An introduction to environments and vegetation. In: B.B. Amos & F.R. Gehlbach, eds. Edwards Plateau vegetation: Plant ecological studies in central Texas. Baylor University Press, Waco, Texas, U.S.A. Pp. 1–15.
- ROBINSON, H.W. and C.T. McALLISTER. 2012. The endemic biota of Oklahoma. Proc. Okla. Acad. Sci. 92:21–28.
- ROHRER, W.L. 1997. Biosystematic study of the rare plant *Paronychia virginica* Sprengel (Caryophyllaceae) employing morphometric and allozyme analyses. MS thesis. Virginia Polytechnic Institute and State University, Blacksburg, Virginia, U.S.A.
- ROGERS, C.M. 1953. The vegetation of the Mesa de Maya region of Colorado, New Mexico, and Oklahoma. Lloydia 16(4):257–290.
- ROGERS, C.M. 1954. Some botanical studies in the Black Mesa region of Oklahoma. Rhodora 56:205–212.
- ROLFSMEIER, S.B. & G. STEINAUER. 2010. Terrestrial ecological systems and natural communities of Nebraska. Nebraska Natural Heritage Program, Nebraska Game and Parks Commission, Lincoln, Nebraska, U.S.A.
- ROWELL, C.M., JR. 1967. Vascular plants of the Texas Panhandle and South Plains. PhD diss. Oklahoma State University, Stillwater, Oklahoma, U.S.A.
- ROWELL, C.M., JR. 1971. Vascular plants of the playa lakes of the Texas Panhandle and South Plains. Southw. Nat. 15(4):407–417.
- RYDBERG, P.A. 1917. Flora of the Rocky Mountains and adjacent plains. Published by the author, New York, New York, U.S.A.
- RYDBERG, P.A. 1931. A short phytogeography of the prairies and Great Plains of central North America. Brittonia 1:57–66 + plate.
- RYDBERG, P.A. 1932. Flora of the prairies and plains of central North America. New York Botanical Garden, New York, New York, U.S.A.
- SAGHATELYAN, A.A. 2015. Phytogeographical relationships and analysis of the flora of south-central Texas, U.S.A. J. Bot. Res. Inst. Texas 9(1):259–294.
- SCHIEBOUT, M.H., D.L. HAZLETT, & N. SNOW. 2008. Floristic survey of the vascular plants over parts of northeastern New Mexico. J. Bot. Res. Inst. Texas 2(2):1407–1447.
- SCHROEDER, W.A. 1982. Presettlement prairie of Missouri, 2nd ed. Natural History Series No. 2, Missouri Department of Conservation, Jefferson City, Missouri, U.S.A.
- SCHULZ, K.A. & R.B. SHAW. 1992. Status of *Haplopappus fremontii* A. Gray ssp. *monocephalus* (A. Nelson) Hall [Asteraceae] in Colorado. Prairie Naturalist 24(3):143–148.
- SCOTT, G.R. & W.A. COBBAN. 1964. Stratigraphy of the Niobrara Formation at Pueblo, Colorado. U.S. Geological Survey Professional Paper 454-L.
- SHAW, R.B., S.L. ANDERSON, K.A. SCHULTZ, & V.E. DIERSING. 1989a. Plant communities, ecological checklist, and species list for the U.S. Army Piñon Canyon Maneuver Site, Colorado. Range Science Series 37. Department of Range Science, Colorado State University, Fort Collins, Colorado, U.S.A.
- SHAW, R.B., S.L. ANDERSON, K.A. SCHULTZ, & V.E. DIERSING. 1989b. Floral inventory for the U.S. Army Piñon Canyon Maneuver Site, Colorado. Phytologia 67:1–42.
- SHULTZ, L.M. 1993. Patterns of endemism in the Utah flora. In: R. Sivinski & K. Lightfoot, eds. Southwestern rare and endangered plants. New Mexico Department of Forestry and Resources Conservation Division, Misc. Publ. No. 2. Santa Fe, New Mexico, U.S.A. Pp. 249–263.
- SHORTHOUSE, J.D. 2010. Ecoregions of Canada's prairie grasslands. In: J.D. Shorthouse & K. D. Floate, eds. Arthropods of Canadian grasslands, vol. 1: Ecology and interactions in grassland habitats. Pp. 53–81.
- SIMS, P.L. & P.G. RISSEY, 2000. Grasslands. In: M.G. Barbour & W.D. Billings, eds. North American terrestrial vegetation, 2nd ed. Cambridge University Press, New York, U.S.A. and Cambridge, U.K. Pp. 323–356.
- SINGH, J.S., W.K. LAURENROTH, R.K. HEITCHMIDT, & J.L. DODD. 1983. Structural and functional attributes of the vegetation of the mixed prairie of North America. Bot. Rev. 49(1):117–149.
- SINGHURST, J.R., L.L. SANCHEZ, D. FRELS, JR., T.W. SCHWERTNER, M. MITCHELL, S. MOREN, & W.C. HOLMES. 2007. The vascular flora of Mason Mountain Wildlife Management Area, Mason County, Texas. S. E. Naturalist 6(4):683–692
- SVINSKI, R.C. & P.J. KNIGHT. 1996. Narrow endemism in the New Mexico flora. In: J. Maschinski, H.D. Hammond, & L. Holter, eds. Southwestern rare and engendered plants: Proceedings of the second conference; 1995 September 11–14, Flagstaff, Arizona. Gen. Tech. Rep. RM-GTR-283, U.S. Department of Agriculture, Forest Service, Fort Collins, Colorado, U.S.A. Pp. 286–296.
- SMITH, B.A. 2010. Four western Cheilantheid ferns in Oklahoma. Oklahoma Native Pl. Rec. 10:65–70.
- SMITH, E.B. 1981. New combinations in *Croptilon* (Compositae-Astereae). Sida 9(1):59–63.

- SMITH, H.L. 1966. Mosses of the Great Plains and Arkansas River lowlands of Kansas. *Univ. Kansas Sci. Bull.* 46(12):433–474.
- SMITH, J.P. & J.O. SAWYER. 1988. Endemic vascular plants of northwestern California and southwestern Oregon. *Madrono* 35:54–69.
- SMITH, L.M. 2003. *Playas of the Great Plains*. University of Texas Press, Austin, Texas, U.S.A.
- SNOW, N. 1990. The distribution of *Frankenia jamesii* Torr. ex Gray (Frankeniaceae) in the Great Plains. *Prairie Nat.* 22(3):211–214.
- SORRIE, B.A. & A.S. Weakley. 2001. Coastal Plain vascular plant endemics: Phytogeographic patterns. *Castanea* 66(1–2):50–82.
- SPEARING, D. 1991. *Roadside geology of Texas*. Mountain Press Publishing Company, Missoula, Montana, U.S.A.
- SPELLENBERG, R.W. 1993. Species of special concern. In: W.A. Dick-Peddie. *New Mexico vegetation: Past, present, and future*. University of New Mexico Press, Albuquerque, New Mexico, U.S.A. Pp. 179–224.
- SPRINGER, T.L. & R.J. TYRL. 1989. Distribution, habitat, and reproductive biology of *Phlox oklahomensis* Wherry (Polemoniaceae). *Proc. Okla. Acad. Sci.* 69:15–21.
- STEBBINS, G.L. & J. MAJOR. 1965. Endemism and speciation in the California flora. *Ecol. Monogr.* 35:1–35.
- STEPHENS, A.R. & W.M. HOLMES. 1989. *Historical atlas of Texas*. University of Oklahoma Press, Norman, Oklahoma, U.S.A.
- SHURE, D.J. 1999. Granite outcrops of the southeastern United States. In: R.C. Anderson, J.S. Fralish, & J.M. Baskin, eds. *Savannas, barrens, and rock outcrop communities of North America*, Cambridge University Press, New York, U.S.A. and Cambridge, U.K. Pp. 99–118.
- SWADEK, R.K. 2012. *Phemaranthus calcaricus* (Montiaceae) new to Texas. *J. Bot. Inst. Texas* 6(10):303–307.
- SWADEK, R.K. & T.L. BURGESS. 2012. The vascular flora of the North Central Texas Walnut Formation. *J. Bot. Res. Inst. Texas* 6(2):725–752.
- SWINEHART, J. B., V. L. Souders, H. M. DeGrew, and R. F. Diffendal, Jr. 1985. Cenozoic paleogeography of western Nebraska. In: R.M. Flores and S.S. Kaplan, eds. *Cenozoic paleogeography of the west-central United States*. Rocky Mountain Paleogeography Symposium 3, Society of Economic Paleontologists and Mineralogists, Rocky Mountain Section, Denver, Colorado. U.S.A. Pp. 209 – 229.
- TAKHTAJAN, A.L. 1986. *Floristic regions of the world*. University of California Press, Berkeley, California, U.S.A.
- TAYLOR, K.N. & R.J. O'KENNON. 2013. Ecology and distribution of the North Central Texas endemic *Dalea reverchonii* (Fabaceae). *J. Bot. Res. Inst. Texas* 7(1):603–610.
- TAYLOR, K.N. & R.J. O'KENNON. 2014. Expanded distribution of *Gratiola quartermaniae* (Plantaginaceae) in Texas, U.S.A. *J. Bot. Res. Inst. Texas* 8(1):333–337.
- TAYLOR, K.N. & R.J. O'KENNON. 2016. The vascular flora of Enchanted Rock State Natural Area, Llano and Gillespie counties, Texas, U.S.A. *J. Bot. Res. Inst. Texas* 10(1):267–294.
- TAYLOR, K.N., R.J. O'KENNON, & T.F. REHMAN. 2012. Expanded distribution of *Isoetes butleri* (Isoetaceae) in Texas. *J. Bot. Res. Inst. Texas* 6:753–757.
- TERLETZKY, P. A. & O.W. VAN AUKEN. 1996. Comparison of cedar glades and associated woodlands of the southern Edwards Plateau. *Texas J. Sci.* 48(1):55–67.
- TERRELL, E.E. 1986. Taxonomic and nomenclatural notes on *Houstonia nigricans* (Rubiaceae). *Sida* 11:471–481.
- THORNE, J.H., J.H. VIERS, J. PRICE, & D.M. STOMS. 2009. Spatial patterns of endemic plants in California. *Nat. Areas J.* 29(4):344–366.
- THORNE, R.F. 1993. Phytogeography. In: *Flora of North America* Editorial Committee, ed. *Flora of North America north of Mexico*, vol. 1. Oxford University Press, New York, U.S.A. and Oxford, U.K. Pp. 132–153.
- TRANSEAU, E.N. 1935. The prairie peninsula. *Ecology* 16:423–427.
- TROCK, D.K. & R.J. O'KENNON. 2003. A new species of *Packeria* (Asteraceae: Senecioneae) from the Edwards Plateau of Texas. *Sida* 20(3):945–951.
- TURNER, B.L. 1959. *The legumes of Texas*. University of Texas Press, Austin, Texas, U.S.A.
- TURNER, B.L. 1993. Texas species of *Mirabilis* (Nyctaginaceae) *Phytologia* 75(6):432–451.
- TURNER, B.L. 1995. Taxonomic overview of *Hedyotis nigricans* (Rubiaceae) and closely allied taxa. *Phytologia* 79(1):12–21.
- TURNER, B.L. 1997. Rebuttal to Terrell's taxonomic notes of Turner's treatment of Texan and Mexican *Hedyotis*. *Phytologia* 82:82–85.
- TURNER, B.L. 2006. *Dalea austrotexana* (Fabaceae), a new species from southernmost Texas. *Phytologia* 88(3):288–293.
- TURNER, B.L. 2009. *Iva corbinii* (Asteraceae): A remarkable new species from Travis County, Texas. *Lundellia* 12:5–7.
- TURNER, B.L. 2012a. A new species of *Cardamine* (Brassicaceae) from south-central Texas. *Phytoneuron* 2012-49:1–5.
- TURNER, B.L. 2012b. Taxonomy of *Euclide bartonioides* (Loaseae) complex in Texas. *Phytologia* 94(3):305–309.

- TURNER, B.L. 2014 Taxonomic overview of *Eustoma* (Gentianaceae). *Phytologia* 96(1):7–11.
- TURNER, B.L. & M.J. MOORE. 2014. *Oenothera gayleana* (*Oenothera* sect. *Calylophus*, Onagraceae), a new gypsophile from Texas, New Mexico, and Oklahoma. *Phytologia* 96(3):200–206.
- TURNER, B.L., H. NICHOLS, G.C. DENNY, & O. DORON. 2003. Atlas of the vascular plants of Texas. *Sida, Bot. Misc.* 24.
- TURNER, M.W. 2003. A new species of *Brazoria* (Lamiaceae) from the Central Mineral Region of Texas. *Sida* 20(4):1565–1571.
- VAN AUKEN, O.W. 2000. Characteristics of intercanopy bare patches in *Juniperus* woodlands of the southern Edwards Plateau, Texas. *Southw. Naturalist* 45(2):95–110.
- WALTERS, T.W. & R. WYATT. 1982. The vascular flora of granite outcrops in the Central Mineral Region of Texas. *Bull. Torrey Bot. Club* 109(3):344–364.
- WARE, S. 2002. Rock outcrop communities (glades) in the Ozarks: A synthesis. *Southw. Naturalist* 47(4):585–597.
- WEAKLEY, A.S., J.C. LUDWIG, & J.F. TOWNSEND. 2012. *Flora of Virginia*. Botanical Research Institute of Texas, Fort Worth, Texas, U.S.A.
- WEAVER, J.E. 1954. *North American prairie*. Johnsen Publishing Company, Lincoln, Nebraska, U.S.A.
- WEBER, W.A. & R.C. WITTMANN. 1992. *Catalog of the Colorado flora: A biodiversity baseline*. University Press of Colorado, Boulder, Colorado, U.S.A.
- WEBER, W.A. & R.C. WITTMANN. 2012. *Colorado flora, eastern slope: A field guide to the vascular plants*, 4th ed. University Press of Colorado, Boulder, Colorado, U.S.A.
- WELLS, P.V. 1970. Historical factors controlling vegetation patterns and floristic distributions in the Central Plains region of North America. In: W. Dort, Jr. & J. K. Jones, eds. *Pleistocene and Recent environments of the central Great Plains*, University of Kansas Department of Geology Special Publication No. 3, University Press of Kansas, Lawrence, Kansas, U.S.A. Pp. 211–221.
- WEST, N.E. 1999. Juniper-pinon savannas and woodlands of western North America. In: R.C. Anderson, J.S. Fralish, & J.M. Baskin, eds. *Savannas, barrens, and rock outcrop communities of North America*, Cambridge University Press, New York, U.S.A. and Cambridge, U.K. Pp. 288–308.
- WHITEHOUSE, E. 1933. Plant succession on central Texas granite. *Ecology* 14(4):391–405.
- WILL-WOLF, S. & F. STEARNS. 1999. Dry soil oak savanna in the Great Lakes region. In: R.C. Anderson, J.S. Fralish, & J.M. Baskin, eds. *Savannas, barrens, and rock outcrop communities of North America*, Cambridge University Press, New York, U.S.A. and Cambridge, U.K. Pp. 135–154.
- WORLD WILDLIFE FUND. 2016. *Plowprint annual report 2016*. World Wildlife Fund Northern Great Plains Program, Bozeman, Montana, U.S.A.
- WRIGHT, R.A. 2001. The vegetation of Palo Duro Canyon. In: D.F. Guy, ed. *The story of Palo Duro Canyon*. Texas Tech University Press, Lubbock, Texas, U.S.A. Pp. 87–116.
- YATSKIEVYCH, G. 1999. *Steyermark's flora of Missouri*, 2nd ed., vol. 1. Missouri Department of Conservation and Missouri Botanical Garden, Jefferson City and St. Louis, U.S.A.
- YATSKIEVYCH, G. 2006. *Steyermark's flora of Missouri*, 2nd ed., vol. 2. Missouri Botanical Garden Press, St. Louis, U.S.A.
- YATSKIEVYCH, G. 2013. *Steyermark's flora of Missouri*, 2nd ed., vol. 3. Missouri Botanical Garden Press, St. Louis, U.S.A.
- ZOLLNER, D., M.H. MACROBERTS, B.R. MACROBERTS, & D. LADD. 2005. Endemic vascular plants of the Interior Highlands, U.S.A. *Sida* 21:1781–1791.